

# Universo Primitivo 2019-2020 (1º Semestre)

Mestrado em Física - Astronomia

## Chapter 1

### 1. The observed Universe

- Foreword: The Olber's paradox;
- The Universe at different scales;
- Observational Cosmology: empirical facts and the hot Big-Bang theory
  - Cosmic Expansion: The Hubble law;
  - The abundancies of the light elements;
  - The existence of a Cosmic Background Radiation;
  - The isotropy of distant objects;
  - The existent of dark matter;
  - The accelerated expansion of the Universe

# Foreword: The Olber's paradox and the present view of the Universe


## Foreword: Why is the sky dark at night?



Heinrich Olbers  
(1758-1840)



# Foreword: Why is the sky dark at night?



Heinrich Olbers  
(1758–1840)

**Olbers' paradox** (1826) : argues that “the **darkness of the sky** at night **conflicts with the concept of an infinite and eternal static universe**”, with stars distributed uniformly.

$$\text{light} = \int_{r_0}^{\infty} L(r)N(r) dr$$



# Foreword: Why is the sky dark at night?




Heinrich Olbers  
(1758–1840)

**Olbers' paradox** (1826) : argues that “the **darkness of the sky** at night **conflicts with the concept of an infinite and eternal static universe**”, with stars distributed uniformly.

Oblers paradox in action.

Exercise: prove why this happens




# Foreword: Why is the sky dark at night?



Heinrich Olber's  
(1758–1840)


Some possible explanations:

1. Too much dust absorbs light from distant stars.
2. The number of stars in the Universe is finite.
3. The distribution of stars is not uniform.
4. The Universe is expanding. Light from distant stars are dimmed (redshifted) into obscurity.
5. The observed Universe has a finite age. Distant light hasn't even reached us yet.




The Universe at different  
scales

Stars...



Sun: 8 light minutes away  
 $\alpha$ -Centauri 4,25 light years




Gaseous Nebulae...




... the birthplaces of stars ...

Distance to the Eagle Nebulae 7000 light years


The galaxy...





A home for billions of stars...



# Milky Way




# Groups and Clusters of Galaxies...




Distance to Coma cluster 321 million ly – 99 Mpc

# Our place in the Universe...



Credits: American Natural History Museum; gently provided by Miguel de Avillez U. Évora

# LSS: “The fingers of God”...



*Credit: M. Blanton and the Sloan Digital Sky Survey*

Further away and back in time...  
First stars forming period

Distance: 12.8 – 13.4 billion years back in time



CMB...

## The edge of the visible Universe

Distance: ~13.8 billion years back in time

Fig. credits: NASA / WMAP Science Team


### Planck Surveyor: looking back to the dawn of time



BR-275

Looking back  
to the dawn of time

PLANCK



European Space Agency  
Agence spatiale européenne

**Project:** ESA lead mission to observe the temperature and polarization anisotropies of the Cosmic Microwave Background (CMB) radiation with unprecedented precision.

**Total Cost:** about €700 million (€1 / person in EU)

#### Mission timeline:

**Launch:** 14 May 2009

**Operational orbit at L2:** July 2009

**Nominal science phase:** end of January 2011

**Extended mission:** Shut down date: 19 Oct. 2013

#### Payload:

**Telescope:** 1.5 m projected apertures


**Low Frequency Instrument (LFI):** array of 22 tuned radio receivers operating at 30, 44 and 70 GHz.

**High Frequency Instrument (HFI):** array of 52 bolometers operating at 100, 143, 217, 353, 545, and 857 GHz.

Fig. credits: ESA

# Planck CMB observations


2009-2013: Planck satellite observes the CMB sky with unprecedented angular resolution and sensitivity.



Animation credits: ESA and the Planck collaboration; Cluster map by Douspis, Hurier, Aghanim 2013

# Planck CMB observations

2009-2013: Planck satellite observes the CMB sky with unprecedented angular resolution and sensitivity.




Animation & Fig. credits: ESA and the Planck collaboration

# Galaxy surveys: 3D mapping of the Universe...

SDSS: aims at ~25% of the sky; ~100 million objects

## ESA Cosmology missions




ESA/NASA Planck Surveyor:  
2009 - 2013

ESA Euclid:  
2021 - 2027



# Euclid mission (ESA): Galaxy Surveys from space (launch 2022)

Portuguese official participation lead by the cosmology group @ IA – CAUP/FCUL





Euclid: ~ 2000 milions of galaxies

Fig. credits: ESA - C. Carreau.

## Euclid survey planning

Portuguese official contribution is carried out at IA/FCUL:

- J. Dínis
- I. Tereno
- C. S. Cavalho
- A. da Silva



From: Euclid Preparation I. The Euclid Reference Survey: status at the Preliminary Design Review, submitted to the ECEB, 2019

# Observational cosmology: empirical facts about the Universe



## 1. The Universe is expanding



Edwin Hubble

1924: Edwin Hubble ends debate on the nature of nebulae being galactic objects

1929: reports a linear relation between relative radial velocity and distance:  $v = Hd$




From:

# 1. The Universe is expanding




Edwin Hubble



1924: Edwin Hubble ends debate on the nature of nebulae being galactic objects

1929: reports a linear relation between relative radial velocity and distance:  $v = Hd$


At a given time



$$d(t) = a(t)d_0$$

From:

At a later time



# 1. The Universe is expanding




Edwin Hubble



1924: Edwin Hubble ends debate on the nature of nebulae being galactic objects

1929: reports a linear relation between relative radial velocity and distance:  $v = Hd$

Time evolution of an expanding spherical surface



$$d(t) = a(t)d_0$$

From:

# The basic idea behind the Big-Bang theory

- If the universe is expanding and matter-energy is conserved during the expansion then the universe had to be smaller, denser and hotter in the past!
- If so, the Universe must have evolved from a state where matter and radiation form a **ultra dense and hot ionized plasma** of fundamental particles
- As the universe expands and cools down:
  - interactions between the plasma components become less frequent;
  - different particle species should decouple from the plasma;
  - eventually the universe becomes neutral and transparent to radiation



## According to the Big-Bang theory, in the early instants...

*“ the Universe was a extremely hot and dense plasma, like a ‘torrid bright fog’ ...*

*... radiation was trapped in this plasma through collisions with other plasma particles*

*... as the universe expands, the plasma temperature drops, atomic nuclei form, and capture the free electrons in the plasma. When the number of free electrons is too small, radiation no longer interacts with the plasma and propagates freely, giving rise to the Cosmic Microwave Background and neutral matter”*

## 2. The abundance of light nuclei




Herman, Gamow, Alpher

The relative abundance of light elements can not be explained by stellar nucleosynthesis

1948: Alpher & Gamow computed the abundance of light elements in the context of the **Big Bang** theory

Light elements were produced at low temperatures ( $< 10^9$  K and high densities) during several tens of minutes



From:



## 3. Cosmic Microwave Background



Penzias & Wilson


1965: Penzias & Wilson serendipitously discovered a uniform radiation (“excess”) across the sky.

This was the cosmic microwave background radiation predicted by Gamow and Alpher in 1948






# CMB: the last scattering surface



33

Reprinted from: <http://physicsworld.com/cws/article/indepth/2014/jan/09/planck-perspectives>

## 3. Cosmic Microwave Background



John Mather &  
Geroge Smoot

1991: High precision measurement of CMB temperature by COBE and 1st detection of temperature fluctuations (Mather & Smoot)

2001: State of the art measurements of  $dT/T \sim 1e-5$  temperature fluctuations by WMAP





Fig. credits: NASA / WMAP Science Team

# 4. Isotropy of distant objects

On Large Scales the Universe...  
... appears to be ISOTROPIC

CMB  
 $T=2.725\text{ K}$

+

The Cosmological Principle (Milne, Einstein):  
"all places in the Universe are alike" – Einstein

The Universe Homogeneous and Isotropic on large scales

The APM Galaxy Survey  
Maddox et al

Distant Objects in the Hubble Ultra Deep Field

NASA, ESA, R. Windhorst (Arizona State University)  
and H. Yan (Spitzer Science Center, Caltech)

# 4. Isotropy of distant objects





Fig. 3.1 A point distribution, statistically isotropic around every point (left) and around a unique point ( $P$ ) (right). In the second version,  $P$  and  $Q$  are not equivalent. The cosmological principle excludes such kinds of solutions, which would assume that we lie in a special place in the Universe. From Ref. [1] of the introduction.


The APM Galaxy Survey  
Maddox et al

NASA, ESA, R. Windhorst (Arizona State University)  
and H. Yan (Spitzer Science Center, Caltech)

# 5. The existence of Dark Matter



From:




Jan Oort

1927: Jan Oort studies the **rotation of stars in our galaxy** and infers that their rotation is not consistent with Keplerian motion.

$$v_{circ} = \sqrt{\frac{GM(R)}{r}}$$

# 5. The existence of Dark Matter



Circular motion:


$$v_{circ} = \sqrt{\frac{GM(R)}{r}}$$

If the whole mass is mostly at the centre:  $v_{cir}^2 \sim 1/r$

Oort's constants:

$$A \equiv -\frac{1}{2} \left[ \frac{dV_c}{dR} \Big|_{R_0} - \frac{V_{c,0}}{R_0} \right]$$

$$B \equiv -\frac{1}{2} \left[ \frac{dV_c}{dR} \Big|_{R_0} + \frac{V_{c,0}}{R_0} \right]$$




Observations vs Keplerian motion:

- **Kepler. motion:**  $(A-B)/(A+B) = 2$
- **Observations :**  $(A-B)/(A+B) = 5$

-Mass is not concentrated at the centre  
-Non-luminous mass is required

## 5. The existence of Dark Matter



B: Observations

A: theoretical expectations

1980: Vera Rubin and others also find that **stars rotate too fast in the outskirts of spiral galaxies** to remain bound assuming that gravity is produced only by visible matter.

## 5. The existence of Dark Matter





Fritz Zwicky

1936: Fritz Zwicky applied the **Virial theorem to the velocities of galaxies in the Coma cluster** and finds very high  $\Upsilon = M/L$  for them to remain bound ( $\Upsilon_{\text{coma}}/\Upsilon_{\text{sun}} \sim 500 \gg 2 - 10$  for galaxies).


- **Virial theorem (for gravitationally relaxed systems):**  $2\bar{E}_k + \bar{E}_p = 0$
- **Mass from the virial theorem:**  $M_V = \langle v^2 \rangle \langle R \rangle / G$
- **Visible (luminous) Mass:**  $M_L = N_g R_{ML} L_g$   
( $R_{ML}$  – typical mass to light ratio of galaxies;  
 $N_g, L_g$  number and luminosity of individual galaxies)

# 5. The existence of Dark Matter

lensing effects:




$$G_{\mu\nu} = 8\pi G T_{\mu\nu}$$



From:

# 5. The existence of Dark Matter

lensing effects:




Strong lensing

From:

# 5. The existence of Dark Matter

lensing effects: **strong lensing**

**Einstein Rings**




Galaxy-galaxy lensing



Double Einstein Ring SDSSJ0946+1006




Hubble Space Telescope • ACS/WFC



NASA, ESA, R. Gavazzi and T. Treu (University of California, Santa Barbara), and the SLACS Team

STScI-PRC08-04

# 6. Cosmic expansion is accelerating




1998: S. Perlmutter and the supernova Cosmology project found first evidence for the accelerated expansion of the Universe.

assuming supernovae are standard candles, they appear further away (**green arrow**) than predicted by non-accelerating expansion models (**yellow arrow**).


$$d(t) = a(t) d_0$$

with  $\ddot{a}(t) > 0$

No Accelerated expansion






Accelerated expansion




From:

# 6. Cosmic expansion is accelerating




Cosmological redshift:

$$\begin{aligned}
 z &= \frac{E - E_0}{E_0} = \\
 &= \frac{\nu}{\nu_0} - 1 = \\
 &= \frac{\lambda_0}{\lambda} - 1 = \\
 &= \frac{a_0}{a} - 1
 \end{aligned}$$



# How Cosmological structure forms and evolves?



Observations indicate that

- ❑ on small scales the universe is NOT homogeneous and isotropic
- ❑ On large cosmological scales the Universe doesn't show indications of strong anisotropies. Together with the cosmological principle this implies the universe is highly homogeneous and isotropic
- ❑ However it shows small anisotropies in the CMB.

# How structure forms and evolves?

Structures should grow due to  
Gravitational Instability (Jeans, Liefshitz,...)

$$\mathbf{g}(\mathbf{r}, t) = -\frac{1}{a} \nabla \phi = \frac{3\Omega H^2}{8\pi} \int d\mathbf{x}' \delta(\mathbf{x}', t) \frac{(\mathbf{x}' - \mathbf{x})}{|\mathbf{x}' - \mathbf{x}|^3}$$

Density fluctuations: t=13.7 billion years

$$\mathbf{g}(\mathbf{r}, t) = -\frac{1}{a} \nabla \phi = \frac{3\Omega H^2}{8\pi} \int d\mathbf{x}' \delta(\mathbf{x}', t) \frac{(\mathbf{x}' - \mathbf{x})}{|\mathbf{x}' - \mathbf{x}|^3}$$



## Density fluctuations: t=13.5 billion years

500 Mpc/h

$$\mathbf{g}(\mathbf{r}, t) = -\frac{1}{a} \nabla \phi = \frac{3\Omega H^2}{8\pi} \int d\mathbf{x}' \delta(\mathbf{x}', t) \frac{(\mathbf{x}' - \mathbf{x})}{|\mathbf{x}' - \mathbf{x}|^3}$$

Millennium simulation, Springel et al.

## Density fluctuations: t=12.7 billion years

500 Mpc/h

$$\mathbf{g}(\mathbf{r}, t) = -\frac{1}{a} \nabla \phi = \frac{3\Omega H^2}{8\pi} \int d\mathbf{x}' \delta(\mathbf{x}', t) \frac{(\mathbf{x}' - \mathbf{x})}{|\mathbf{x}' - \mathbf{x}|^3}$$

Millennium simulation, Springel et al.

## Density fluctuations: t=9 billion years

500 Mpc/h

$$\mathbf{g}(\mathbf{r}, t) = -\frac{1}{a} \nabla \phi = \frac{3\Omega H^2}{8\pi} \int d\mathbf{x}' \delta(\mathbf{x}', t) \frac{(\mathbf{x}' - \mathbf{x})}{|\mathbf{x}' - \mathbf{x}|^3}$$

Millennium simulation, Springel et al.

## Density fluctuations: t=1 billion years

500 Mpc/h

$$\mathbf{g}(\mathbf{r}, t) = -\frac{1}{a} \nabla \phi = \frac{3\Omega H^2}{8\pi} \int d\mathbf{x}' \delta(\mathbf{x}', t) \frac{(\mathbf{x}' - \mathbf{x})}{|\mathbf{x}' - \mathbf{x}|^3}$$

Millennium simulation, Springel et al.

# Cosmological structure formation

$z = 20.0$

50 Mpc/h



# Large Scale Structure (LSS)

1 Gpc/h




Millennium Simulation

10,077,696,000 particles

( $z = 0$ )

# Large Scale Structure (LSS)



# The history of the Universe:

