

Fundamentos de Biologia Molecular

Curso de Licenciatura em Biologia
2º Ano, 1º Semestre
Ano Letivo 2018/2019

Componente Teórico-Prática



Ciências
ULisboa

Faculdade
de Ciências
da Universidade
de Lisboa


Docente Responsável: Rita Zilhão
Docente TPs: Andreia Figueiredo

TP2: Polymerase Chain Reaction (PCR)

- DNA structure
- PCR discovery
- Nobel prizes
- PCR
- PCR components
- Setting up a PCR reaction
- Practical applications



TP2: Polymerase Chain Reaction (PCR)

The structure of DNA




Double Helix

Complementary Base Pairing



Antiparallel Strands



Unzipping

TP2: Polymerase Chain Reaction (PCR)

The structure of DNA

How do we identify and detect a specific sequence in a genome?

- There are a LOT of other sequences in a genome that we're not interested in detecting. (SPECIFICITY)
- The amount of DNA in samples we're interested in is VERY small. (AMPLIFICATION)

How do we identify and detect a specific sequence in a genome?

- Pine: 68 billion bp
- Corn: 5.0 billion bp
- Soybean: 1.1 billion bp
- Human: 3.4 billion bp
- Housefly: 900 million bp
- Rice: 400 million bp
- E. coli: 4.6 million bp
- HIV: 9.7 thousand bp



TP2: Polymerase Chain Reaction (PCR)

Specificity

- The human genome is 3.4 B bp
- If the bases were written in standard 10-point type, on a tape measure...
...The tape would stretch for 8635.73 Km!



Identifying a 500bp sequence in a genome would be like finding a section of this tape measuring only 1.21 meters long!



TP2: Polymerase Chain Reaction (PCR)

Detection

How many molecules do we need to be able to see them?

- We need a minimum of 10 ng of fluorescent stained DNA to detect on agarose gel

URI Genomics & Sequencing Center

Calculator for determining the number of copies of a template


enter amount of DNA (ng):

enter length of template (bp):


Calculate number of copies!

number of copies : $\times 10^{\wedge}10$

In other words, to “see” a single “gene”, the DNA in a sample of 100 cells would have to be multiplied 180 million times!!!!




TP2: Polymerase Chain Reaction (PCR)



PCR is widely used to amplify target DNA regions of known sequences within a heterogeneous collection of DNA sequences

- Taq DNA polymerase isolation from *Thermus aquaticus* by Brock & Freeze - 1969
- Kary Mullis, PCR invention - 1988

Habitat of
Thermus aquaticus



Structure of Taq Polymerase

TP2: Polymerase Chain Reaction (PCR)



Kary B. Mullis

- The inventor of the DNA synthesis process known as the **Polymerase Chain Reaction (PCR)**. The process is an invaluable tool to today's molecular biologists and biotechnology corporations.
- Mullis, born in Lenoir, North Carolina, attended the University of Georgia Tech for his undergraduate work in chemistry, and then obtained a Ph. D. in biochemistry from Cal Berkeley.
- In 1983, working for Cetus Corporation, Mullis developed the Polymerase Chain Reaction, a technique for the rapid synthesis of a DNA sequence. ***The simple process involved heating a vial containing the DNA fragment to split the two strands of the DNA molecule, adding oligonucleotide primers to bring about reproduction, and finally using polymerase to replicate the DNA strands. Each cycle doubles the amount of DNA, so multiple cycles increase the amount of DNA exponentially, creating huge numbers of copies of the DNA fragment.***

TP2: Polymerase Chain Reaction (PCR)



Michael Smith

- Smith first conceived of ***site-directed mutagenesis*** in the early 1970s and devoted several years to working out the details of the technique. The method provided researchers with a new way to study protein function.
- A protein is a compound made up of strings of amino acids that fold into a three-dimensional structure, and the protein's structure determines its function. Instructions for the amino-acid sequence of a protein are contained in its gene, namely, in the sequence of DNA subunits, called nucleotides, that make up that gene. The amino-acid sequence of a protein, and hence its function, can be modified by inducing mutations in the nucleotide sequence of its gene. Once an altered protein has been produced, its structure and function can be compared to those of the natural protein.
- Smith developed site-directed mutagenesis, a technique that can be used to modify nucleotide sequences at specific, desired locations within a gene. ***This has made it possible for researchers to determine the role each amino acid plays in protein structure and function.*** Aside from its value to basic research, site-directed mutagenesis has many applications in medicine, agriculture, and industry.

TP2: Polymerase Chain Reaction (PCR)



Kary B. Mullis



Michael Smith

The Nobel Prize in Chemistry 1993 was awarded "***for contributions to the developments of methods within DNA-based chemistry***" jointly with one half to Kary B. Mullis "for his invention of the polymerase chain reaction (PCR) method" and with one half to Michael Smith "***for his fundamental contributions to the establishment of oligonucleotide-based, site-directed mutagenesis and its development for protein studies***".

TP2: Polymerase Chain Reaction (PCR)


PCR

- Developed by Kary Mullis in 1985,
- PCR results in the selective amplification of a chosen region of a DNA molecule,
- The borders of the amplified region have to be known and two short oligonucleotides have to be synthesized,
- These oligonucleotides, which act as ***primers*** for the DNA synthesis, delimit the region to be amplified.




Exponential copies of template DNA


TP2: Polymerase Chain Reaction (PCR)



Parental strands



Two oligonucleotides (**primers**) that hybridize in opposite strands and flank the target




Repetitive cycles:

Template denaturation


Primer annealing

Extension


TP2: Polymerase Chain Reaction (PCR)




TP2: Polymerase Chain Reaction (PCR)



TP2: Polymerase Chain Reaction (PCR)



TP2: Polymerase Chain Reaction (PCR)



$$2^1 = 2$$

$$2^2 = 4$$

$$2^3 = 8$$

In each cycle:
D- denaturation
A- annealing
E- extension

No. of copies/each initial molecule = 2^n
n, no. of cycles

TP2: Polymerase Chain Reaction (PCR)




Table 18.5

Number of copies of DNA fragment in PCR amplification

Number of PCR Cycles (<i>n</i>)	Number of Double-Stranded Copies of Original DNA (2 ^{<i>n</i>})
0	1
1	2
2	4
3	8
4	16
5	32
6	64
7	128
8	256
9	512
10	1,024
20	1,048,576
30	1,073,741,824

n, no. of cycles

TP2: Polymerase Chain Reaction (PCR)

PCR






Fig. 1. The PCR temperature cycle: (1) the temperature is raised to about 95 °C to melt the double stranded DNA, (2) the temperature is lowered to let primers anneal, (3) the temperature is set to 72 °C to let the polymerase extend the primers.

- ★ melting temperature (T_m)- dependent
- ★ Size- and polymerase- dependent

TP2: Polymerase Chain Reaction (PCR)

PCR reaction

- Water
- Buffer
- DNA template
- Primers
- Nucleotides
- Mg^{2+} ions
- DNA Polymerase



TP2: Polymerase Chain Reaction (PCR)

PCR reaction components

- **Water**
- Buffer
- DNA template
- Primers
- Nucleotides
- Mg^{2+} ions
- DNA Polymerase

- **Water**
 - The medium for all other components.



TP2: Polymerase Chain Reaction (PCR)

PCR reaction components

- Water
- **Buffer**
- DNA template
- Primers
- Nucleotides
- Mg^{2+} ions
- DNA Polymerase

- **Buffer**

- Stabilizes the DNA polymerase, DNA, and nucleotides
- 500 mM KCl
- 100 mM Tris-HCl, pH 8.3
- Triton X-100 or Tween




TP2: Polymerase Chain Reaction (PCR)

PCR reaction components

- Water
- Buffer
- **DNA template**
- Primers
- Nucleotides
- Mg^{2+} ions
- DNA Polymerase

- **DNA template**
 - Contains region to be amplified
 - Any DNA desired
 - Purity not required
 - Should be free of polymerase inhibitors



TP2: Polymerase Chain Reaction (PCR)

PCR reaction components

- Water
- Buffer
- DNA template
- **Primers**
- Nucleotides
- Mg²⁺ ions
- DNA Polymerase


- **Primers**

- Specific for ends of amplified region
- Forward and Reverse
- Annealing temps should be known
 - Depends on primer length, GC content, etc.
- Length 15-30 nt
- Conc 0.1 – 1.0 μM (pMol/ μl)



TP2: Polymerase Chain Reaction (PCR)

Choosing primer sequences



TP2: Polymerase Chain Reaction (PCR)

Primer design tips

- Typical PCR primers are anything between 18-28 nucleotides in length
- The G+C composition should ideally be similar to that of the desired amplicon and should in general be between 40-60%
- The calculated T_m for a primer pair should be balanced
- Rule of thumb: $T_m = 2(A+T) + 4(G+C)$ and $-1.5\text{ }^\circ\text{C}$ for every mismatch
- A T_m 55°C - 72°C is desired (62 - 65°C is best)
- Check for complementarity in 3' ends of primer pairs - this leads to primer - dimer artifacts
- Avoid any significant secondary structure within primers i.e. internal palindromic sequences
- Runs of 3 or more C's and G's at 3' ends promote mispriming in G/C rich regions
- Avoid an A and especially a T at the 3' end of a primer
- Avoid any potential mismatches in the 3' end of primers (pay attention when designing degenerate primers)

TP2: Polymerase Chain Reaction (PCR)

Primer design using Primer3

Website: <http://bioinfo.ut.ee/primer3-0.4.0/>

Place here query sequence for which primers will be designed

Min-Max size of generated amplicon (also counting with intermediate values)

e.g. – 200-500

Amplicons with size ranging from 200 bp to 500 bp

Min, Max and Opt size/melting temperature and GC content of primers

Primer3 (v. 0.4.0) Pick primers from a DNA sequence.

Checks for mispriming in template.	disclaimer	Primer3 Home
Primer3plus interface	cautions	FAQ/WIKI

There is a newer version of Primer3 available at <http://primer3.ut.ee>

Paste source sequence below (5'→3', string of ACGTNacgtm -- other letters treated as N -- numbers and blanks ignored). FASTA format ok. Please N-out undesirable sequence (vector, ALUs, LINEs, etc.) or use a [Mispriming Library \(repeat library\)](#): NONE

Pick left primer, or use left primer below: Pick hybridization probe (internal oligo), or use oligo below: Pick right primer, or use right primer below (5' to 3' on opposite strand):

[Sequence Id:](#) A string to identify your output.

[Targets:](#) E.g. 50,2 requires primers to surround the 2 bases at positions 50 and 51. Or mark the [source sequence](#) with [and]: e.g. ...ATCT[CCCC]TCAT. means that primers must flank the central CCCC.

[Excluded Regions:](#) E.g. 401,7 68,3 forbids selection of primers in the 7 bases starting at 401 and the 3 bases at 68. Or mark the [source sequence](#) with < and >: e.g. ATCT CCCC TCAT <68,3> forbids primers in the central CCCC.

[Product Size Ranges](#) 150-250 100-300 301-400 401-500 501-600 601-700 701-850 851-1000

Number To Return 5	Max 3' Stability 9.0
Max Repeat Mispriming 12.00	Pair Max Repeat Mispriming 24.00
Max Template Mispriming 12.00	Pair Max Template Mispriming 24.00

General Primer Picking Conditions

Primer Size Min: 18 Opt: 20 Max: 27	
Primer Tm Min: 57.0 Opt: 60.0 Max: 63.0	Max Tm Difference: 100.0 Table of thermodynamic parameters: Breslauer et al. 1986 ▼
Product Tm Min: Opt: Max:	
Primer GC% Min: 20.0 Opt: Max: 80.0	

Max Self Complementarity: 8.00	Max 3' Self Complementarity: 3.00
Max #N's: 0	Max Poly-X: 5
Inside Target Penalty:	Outside Target Penalty: 0 Note: you can set Inside Target Penalty to allow primers inside a target.
First Base Index: 1	CG Clamp: 0
Concentration of monovalent cations: 50.0	Salt correction formula: Schildkraut and Lifson 1965 ▼
Concentration of divalent cations 0.0	Concentration of dNTPs 0.0
Annealing Oligo Concentration: 50.0 (Not the concentration of oligos in the reaction mix but of those annealing to template.)	

[Liberal Base](#) [Show Debugging Info](#) Do not treat ambiguity codes in libraries as consensus [Lowercase masking](#)

TP2: Polymerase Chain Reaction (PCR)

5' ... CCTGACAGCTACGATACGAGGGAT ATGCTGATGACAGTAGATCTAGTA ... 3'
3' ... GGACTGTCGATGCTATGCTCCCTA TACGACTACTGTCATCTAGATCAT ... 5'



5' ... CCTGACAGCTACGATACGAGGGAT ATGCTGATGACAGTAGATCTAGTA ... 3'
3' ... GGACTGTCGATGCTATGCTCCCTA TACGACTACTGTCATCTAGATCAT ... 5'



Heat

5' ... CCTGACAGCTACGATACGAGGGAT ATGCTGATGACAGTAGATCTAGTA ... 3'
5' GACAGCTACGATACGAG
3' ... GGACTGTCGATGCTATGCTCCCTA TACGACTACTGTCATCTAGATCAT ... 5'



5' ... CCTGACAGCTACGATACGAGGGAT ATGCTGATGACAGTAGATCTAGTA ... 3'
5' GACAGCTACGATACGAGGGAT
3' ... GGACTGTCGATGCTATGCTCCCTA TACGACTACTGTCATCTAGATCAT ... 5'

TP2: Polymerase Chain Reaction (PCR)

Primer storage

- Oligonucleotides are chemically stable. Left dry, they should be good for years. Once hydrated, they are susceptible to degradation by nucleases. If handled correctly, they should be stable for years
- Any DNA oligonucleotide can be degraded by microbial or fingertip nucleases
- Primers should be stored in deionized water or buffered solutions containing EDTA and kept frozen when not in use
- Make a concentrated stock solution in water or TE (10 mM Tris pH 8.0, 1 mM EDTA). A convenient stock could be 100 μ M, stored at -20°C
- Stock solutions should be distributed into several tubes for long term storage so that accidental contamination of a tube will not lead to loss of the entire synthesis


TP2: Polymerase Chain Reaction (PCR)

PCR reaction components

- Water
- Buffer
- DNA template
- Primers
- **Nucleotides (dNTPs)**
- Mg^{2+} ions
- DNA Polymerase


- **Nucleotides (dNTPs)**

- Added to the growing chain
- Activated NTP's
- dATP, dGTP, dCTP, dTTP
- Stored at 10mM, pH 7.0
- Add to 20-200 μ M in assay




TP2: Polymerase Chain Reaction (PCR)

PCR reaction components



Purines



Pyrimidines



dNTP, deoxyribonucleotide triphosphate. Each dNTP is made up of a phosphate group, a deoxyribose sugar and a nitrogenous base. There are four different dNTPs and can be split into two groups: the purines (**A,G**) and the pyrimidines (**T,C**) that connect through hydrogen bridges.

This figure also shows the five common nitrogenous bases found in DNA and RNA on the right.


TP2: Polymerase Chain Reaction (PCR)

PCR reaction components

- Water
- Buffer
- DNA template
- Primers
- Nucleotides
- **Mg²⁺ ions**
- DNA Polymerase

- **Mg²⁺ ions**

- Essential co-factor of DNA polymerase
- Insufficient concentration reduces yield
- Stabilizes the DNA double-helix
- Too much: DNA extra stable, non-specific priming, band smearing
- Used at 0.5 to 3.5 μM in the assay




TP2: Polymerase Chain Reaction (PCR)

PCR reaction components

- Water
- Buffer
- DNA template
- Primers
- Nucleotides
- Mg²⁺ ions
- **DNA Polymerase**

- **DNA Polymerase**
 - The enzyme that does the extension
 - TAQ or similar
 - Heat-stable
 - Approx 1 U / rxn




TP2: Polymerase Chain Reaction (PCR)

Thermostable DNA polymerase (taq DNA polymerase)

When studies of biological organisms in hot springs began in the 1960s, scientists thought that the life of thermophile bacteria could not be sustained in temperatures above about 55 °C.


Thomas D. Brock and Hudson Freeze (Indiana University) reported a new species of thermophile bacterium which they named ***Thermus aquaticus***. The bacterium was first discovered in the Lower Geyser Basin of Yellowstone National Park, near the major geysers and has since been found in similar thermal habitats around the world.



Taq DNA polymerase was isolated from *T. aquaticus* in 1969 by Brock & Freeze 1969


TP2: Polymerase Chain Reaction (PCR)

Structure of DNA Polymerase



TP2: Polymerase Chain Reaction (PCR)


Proofreading



Extension proceeds along the template strand at the 3' end of the newly synthesized strand. When the polymerase recognizes an error, the mismatched base is transferred to the exonuclease active site and the base is excised. The extended strand returns to the polymerase domain, re-anneals to the template strand, and replication continues.

TP2: Polymerase Chain Reaction (PCR)

Proofreading




	KOD (%)	Phusion HF (%)	Pt <i>Taq</i> (%)	Expand HF (%)	FastStart HF (%)	Sequal Prep Long (%)
Overall error rate ^a	0.21	0.11	0.34	0.25	0.23	0.29
Insertions	0.10	0.07	0.14	0.11	0.11	0.11
Deletions	0.06	0.02	0.08	0.07	0.05	0.06
Substitutions	0.01	0.01	0.07	0.04	0.03	0.07
Dots or Dot ^b	0.04	0.01	0.05	0.04	0.04	0.05


TP2: Polymerase Chain Reaction (PCR)

5' – 3' exonuclease activity

DNA Polymerase I 5'→3' Exonuclease activity




DNA polymerase I removes forward (5' to 3') the nucleotides in its way and adds new ones



TP2: Polymerase Chain Reaction (PCR)

Klenow fragment



Klenow fragment (68 kDa)

- 5'-3' polymerase activity
- 3'-5' exonuclease activity

DNA polymerase I (103 kDa)

- 5'-3' polymerase activity
- 5'-3' exonuclease activity
- 3'-5' exonuclease activity

TP2: Polymerase Chain Reaction (PCR)

Example of a PCR reaction

Sterile Water	38.0 μ l
10X PCR Buffer	5.0 μ l
MgCl ₂ (50mM)	2.5 μ l
dNTP's (10mM each)	1.0 μ l
PrimerFWD (25 pmol/ μ l)	1.0 μ l
PrimerREV	1.0 μ l
DNA Polymerase	0.5 μ l
DNA Template	1.0 μ l
Total Volume	50.0 μ l

TP2: Polymerase Chain Reaction (PCR)

Example of a PCR reaction


Sterile Water	38.0 μl
10X PCR Buffer	5.0 μl
MgCl ₂ (50mM)	2.5 μl
dNTP's (10mM each)	1.0 μl
PrimerFWD (25 pmol/ μl)	1.0 μl
PrimerREV	1.0 μl
DNA Polymerase	0.5 μl
DNA Template	1.0 μl
Total Volume	50.0 μl

Mixing Common Reagents Saves Time – **master mix** concept

Component	1X (μ)	20X (μl)
Sterile Water	38.0	760
10X PCR Buffer	5.0	100
MgCl ₂ (50mM)	2.5	50
dNTP's (10mM each)	1.0	20
PrimerFWD (25 pmol/ μl)	1.0	20
PrimerREV	1.0	20
DNA Polymerase	0.5	10
DNA Template	1.0	--
Total Volume	50.0	980



Aliquot
49 μl



TP2: Polymerase Chain Reaction (PCR)

Thermal cycling conditions

Step	Time	Temperature	Comments
1. Denaturation	4 min	94°C	Denaturation of template and primer-dimers
2. Denaturation cycle	0.5 min	94°C	
3. Annealing cycle	0.5 min	55°C	Or 5°C below the lowest primer melting temperature
4. Extension cycle	1 min/kb	72°C	
5. Repeat steps 2–4			Repeat 30 times
6. Final extension	10 min	72°C	
7. End	Infinite	4°C	



TP2: Polymerase Chain Reaction (PCR)

Thermal cycling steps

Denature DNA sample
to separate DNA strands
(94°C, 5 min)

**Initial
DENATURATION**
(Strand separation)

Primers bind to
DNA strands
(50°-65°C, 30 s)

ANNEALING
(Primer hybridization)

Polymerase synthesizes
new DNA strands
(65°-75°C, 2-5 min)

EXTENSION
(DNA synthesis)

X 20-30 cycles

DENATURATION
(Strand separation)


Denature to separate
DNA strands
(94°C, 30 s)

TP2: Polymerase Chain Reaction (PCR)

Visualization of PCR products




- After thermal cycling, tubes are taken out of the PCR machine.
- Contents of tubes are loaded onto an agarose gel.
- DNA is separated by size using an electric field.
- DNA is then stained.
- PCR products are visible as different “bands”.




TP2: Polymerase Chain Reaction (PCR)

Visualization of PCR products



TP2: Polymerase Chain Reaction (PCR)


Visualization of PCR products



TP2: Polymerase Chain Reaction (PCR)

Visualization of PCR products

The final result of the traditional PCR procedure is a gel with a series of bands:





Bands can be compared against each other, and to known size-standards, to determine the presence or absence of a specific amplification product.

TP2: Polymerase Chain Reaction (PCR)

Visualization of PCR products

- End point detection
- Low resolution



Applications

TP2: Polymerase Chain Reaction (PCR)


1. Diagnosis of pathogens	a) PCR b) Nested PCR c) Quantitative PCR d) Multiplex PCR e) Differential on-line and real time PCR
2. Typing genetic markers	a) RFLPs b) AFLPs c) Short tandem repeat polymorphisms
3. DNA template for	a) Genomic mutation screening intron-specific primers flanking exons b) RT-PCR cDNA used as templates for pairs of exon-specific primers to generate overlapping fragments
4. Detecting point mutations	a) Restriction site polymorphisms b) Allele specific amplification
5. cDNA cloning	a) DOP-PCR b) RACE
6. Genomic DNA cloning	New members of a DNA family a) DOP-PCR. Whole genome or subgenomic amplification b) DOP-PCR c) Linker-primed PCR
7. Genome walking	a) Inverse PCR b) Bubble linker (vectored) PCR c) IRE-PCR
8. DNA templates for DNA sequencing	a) ssDNA by asymmetric PCR b) dsDNA for direct sequencing or for cloning followed by sequencing
9. <i>In vitro</i> mutagenesis	a) 5' add-on mutagenesis to create a recombinant PCR product b) Mismatched primers to change a single predetermined nucleotide

TP2: Polymerase Chain Reaction (PCR)

- Amplification of specific DNA fragments to clone, or to isolate a specific clone (gene discovery)
- Distinction between alleles
- Detection of polymorphisms, Molecular Typing, Taxonomy, Forensics, Molecular Diagnosis
- Phylogenetic studies
- Direct isolation of a specific segment of DNA of different origins (genomic, plasmidic etc)
- Preparation of probes
- Tagging of genes by insertion mutations, site-directed mutagenesis
- Cloned DNA molecules are sequenced rapidly by new methods based on PCR

TP2: Polymerase Chain Reaction (PCR)

a)



Transform bacterial with vector
Grow bacterial colonies on agar plate

Isolate DNA from individual colonies

b)



PCR used to amplify the
cloned DNA using primers
for the vector sequences




The cloned DNA
insert is amplified

c)



Analyze PCR products on gel

Amplification by PCR of a specific target region from total genomic DNA for use in cloning




Detection of polymorphisms

-**Tandem repeats** - copies which lie adjacent to each other, either directly or inverted: **Satellite DNA, Minisatellite, Microsatellite**



Tandem byke



(1) Satellite DNA-typically in centromeres and heterochromatin (2) Minisatellite- repeat units of more than 10 base pairs, in many places in the genome, inc. centromeres (3) Microsatellite-repeat units of less than 10 base pairs; incl. telomeres

Detection of polymorphisms



VNTR (Variable Number of Tandem Repeats)

Minisatellites or Microsatellites (STRs or SSRs)

Polymorphism in microsatellites can arise from polymerase slippage

In plants, the most common plant repeat motif is AA/TT, followed by AT/TA and CT/GA.

In mammals the GT/CA motif, being the most abundant dinucleotide repeat, was found to be considerably less frequent in plants.




Detection of polymorphisms

VNTR (Variable Number of Tandem Repeats)


Minisatellites or Microsatellites (STRs or SSRs)

Principle of polymorphism detection (ex. VNTR)



Alleles amplified by PCR are separated in agarose gels

Polymorphism detection Case study





Key

PCR primers

Microsatellite repeats

P Dominant disease allele


$M' - M''''$ Molecular markers



Detection of polymorphisms


VNTR detection by PCR using multiple primers

Multiplex reaction




Detection of polymorphisms

Microsatellites detected by capillary electrophoresis - Fragment Analysis




Fragment analysis detects changes in the length of a specific DNA sequence to indicate the presence or absence of a genetic marker



Phylogenetic studies

Ex. Amplification of 16S rRNA



Other DNA regions frequently used in phylogenetic studies: rRNA

16S

18S

25S/26S

ITS, internal transcribed spacer, between 16S and 23S in bacteria or between 18S and 25S/26S in plants

trnL

matK

rbcL

Cox1

...