

# Index

---

*Note:* Page numbers followed by “f” indicate figures, “t” indicate tables and “b” indicate boxes.

## A

Acenaphthene (Ace), 187–188  
Acenaphthylene (Acy), 187–188  
Adaptation, 105, 115, 117  
Adaptive management (AM), 215  
Adjacency matrix in modeling, 18f, 20–22  
Agricultural products contamination, 166–172  
  Danish agriculture land  
    cadmium balance, 168f  
    lead balance of average, 167f  
  model, 168–171  
    equations, 170t  
    results, 171–172  
  objective, 167  
Air sampling, 187–188  
Aliphatic compounds, 163–164  
Allometric estimation methods, 164  
AM. *See* Adaptive management (AM)  
*Anabaena spirodies*. (*A. spirodies*), 83  
ANNs. *See* Artificial neural networks (ANNs)  
Anthracene (Ant), 187–188  
Antibiotics in wastewater, 173  
Aquatic ecosystem, 66  
Aquatic plants, 185–186  
Aquatic vegetation, 184  
Arrhenius expression, 117  
Artificial neural networks (ANNs), 6–7, 124,  
  124f, 127f  
  MLP, 125–132  
    advantages and disadvantages, 133–134  
    in ecological modeling, 132  
    usage in ecological modeling, 134–138

## B

BaA. *See* Benzo[a]anthracene (BaA)  
Bacillariophytes (Baci), 73–74  
Backpropagation (BP), 124, 127–128  
  algorithm in MLP, 129b  
  learning algorithm, 126–128  
  forward propagation, 126–127  
*Ballamya purificata* (*B. purificata*), 76  
BaP. *See* Benzo[a]pyrene (BaP)

Barrier island vegetation, 239  
BbF. *See* Benzo[b]fluoranthene (BbF)  
BCF. *See* Biological concentration factor (BCF)  
Benefit-cost analysis, 153  
Benzo[a]anthracene (BaA), 187–188  
Benzo[a]pyrene (BaP), 186  
Benzo[b]fluoranthene (BbF), 187–188  
Benzo[ghi]perylene (BghiP), 187–188  
Benzo[k]fluoranthene (BkF), 187–188  
BghiP. *See* Benzo[ghi]perylene (BghiP)  
Bilinear collision, 48  
Bioconcentration factors. *See* Biological concentration factor (BCF)  
Biodegradation, 162, 173  
  rate, 163  
Biodemographic model, 4–5, 5t  
Bioenergetic model, 4–5, 5t  
Biogeochemical ecological models, 145  
Biogeochemical models, 4–7, 5t. *See also* Dynamic population models; Steady state models  
  advantages and properties, 13–16  
  ecological models, 33–37, 34f  
  environmental biogeochemical models, 33–37, 34f, 36t  
  modeling elements, 16–19  
  modeling procedure, 19–33, 23f  
Biological concentration factor (BCF), 154–158,  
  162, 184  
  regression equations for estimation, 162t  
Biological neurons, 124, 124f  
BkF. *See* Benzo[k]fluoranthene (BkF)  
BP. *See* Backpropagation (BP)

## C

Cadmium (Cd), 168  
  balance of average Danish agriculture land, 168f  
  Cd model, 169f  
  concentration in plants and soil, 171f  
  contamination of agricultural products by, 166–172  
CADS. *See* Computer-Aided Design Systems (CADS)  
CalCOFI. *See* California Cooperative Oceanic Fisheries Investigations (CalCOFI)  
Calibration, 24–25, 28

- California Cooperative Oceanic Fisheries Investigations (CalCOFI), 251
- California Current system (CCS), 251  
 climate change impacts on, 250–254  
 evolution of CCS, 252–254, 253f  
 modeling of CCS, 251–252
- Cascadia National Park (CNP), 213  
 FLAM model for, 213–215, 214f
- Causality, 15–18
- CCM. *See* Climate change models (CCM)
- CCS. *See* California Current system (CCS)
- Chlorophytes (Chlo), 73–74
- Chrysene (Chr), 187–188
- Ciprofloxacin, 176
- Cladocerans (Clad), 81–82
- Classical stepwise method, 131–132
- Climate change  
 ADM application for ecological change assessment, 117–121  
 fuzzy effects matrix for, 220t  
 preferred management actions for, 217–220  
 scenarios, 240–241, 241t  
 weighted normalized fuzzy effects matrix, 221t
- Climate change models (CCM), 6–7
- CNP. *See* Cascadia National Park (CNP)
- Coast-wide Reference Monitoring System (CRMS), 234
- Coastal ecosystem modeling in climate change  
 on CCS, 250–254  
 challenges in predicting effects, 254  
 coastal ecosystem processes, 250  
 on coastal ecosystems in Western Pacific, 245–250  
 on Gulf of Mexico Hypoxia, 243–245  
 on Louisiana's coastal wetland plant communities, 230–243  
 tools for assessing effects, 228–229
- COAWST. *See* Coupled Ocean Atmosphere Wave and Sediment Transport modeling system (COAWST)
- Coefficients of variation (CVs), 195, 203
- Competition models, 52–54  
 populations competing for food, 54f
- Computer-Aided Design Systems (CADS), 98
- Conceptual diagram, 18  
 nitrogen cycle in lake, 18f  
 STELLA, 24f
- Consumption/biomass ratio (Q/B ratio), 73–74
- Contribution of variables, 136–138  
 partial derivatives of MLP model, 137f  
 relative contribution of input variables, 137f
- Control functions, 16–18
- Cooperation models, 55–56
- Copepods (Cope), 81–82
- Corbicula fluminea* (*C. fluminea*), 81–82
- Coupled Ocean Atmosphere Wave and Sediment Transport modeling system (COAWST), 248
- Coupled system, 212–215
- Crisp decision rules, 212
- CRMS. *See* Coast-wide Reference Monitoring System (CRMS)
- CVs. *See* Coefficients of variation (CVs)
- Cyanobacteria (Cyan), 73–74
- Cyclic causal approach, 9
- Cycling of matter, 58–61  
 dead matter, 59–60  
 food chain with two trophic levels and cycling of nutrients, 60–61  
 nutrient, 59–60, 59f  
 phytoplankton, 59–60
- ## D
- DahA. *See* Dibenz[a,h]anthracene (DahA)
- Daphnia hyalina* (*D. hyalina*), 81–82
- Daphnia pulex* (*D. pulex*), 81–82
- Darwin's finches, 115–116, 116f
- Darwin's theory, 107–108
- Data preparation, 134
- Data preprocessing, 130–131
- DDT model, 147–148, 148f
- Dead matter, 59–60
- Dibenz[a,h]anthracene (DahA), 187–188
- DIC. *See* Dissolved inorganic carbon (DIC)
- Differential equations, 13–14, 14f, 24, 42
- Dissolved inorganic carbon (DIC), 250–251
- Dust samples, 187–188
- Dynamic equilibrium, 66
- Dynamic model of toxic substance, 147–148
- Dynamic population models. *See also* Biogeochemical models; Steady state models  
 competition models, 52–54  
 cooperation models, 55–56  
 cycling of matter, 58–61  
 first law of population dynamics, 40–43  
 food chains, 56–58, 56f  
 logistic population  
 in periodic environment, 44–45, 46f  
 in random environment, 46–48  
 population model, 39–40  
 prey–predator models, 48–52, 48f, 50f  
 second law of population dynamics, 43–44
- ## E
- Earth's surface modeling, 91. *See also* Ecological modelling  
 EQIs, 92–93  
 FTESM and corollaries, 97–98  
 HASM, 93–97
- Eco-exergy, 107–109, 109f, 111–114, 117  
 $\beta$  values, 112, 113t–114t  
 density, 111  
 information, 111

- Ecological law of thermodynamics (ELT), 112
- Ecological magnification factor (EMF), 162
- Ecological modelling, 13, 33–37, 34f, 103–104
- application, 1–4
    - classification and types, 4–7
    - ecosystem as object for modeling, 9–10
    - environmental problems, 2f
    - model types, 7–9, 8t
  - development, 20
  - MLP, 132
  - MLP usage
    - contribution of variables, 136–138
    - data, 134, 135t
    - data preparation, 134
    - model training, 135, 135f
    - problem, 134
    - results, 135–136, 136f
    - recommendations, 134
- Ecological thermodynamic theory, 66
- Ecopath model, 66–67
- development
    - construction and parameter, 73–74
    - determination of TL, 76
    - evaluation of ecosystem functioning, 74–76
  - for large Chinese lake, 68–85
    - natural system, 68–69
    - study site, 70
    - TLs, 69
  - results and discussion
    - basic model performance, 76–79
    - changes in ecosystem functioning, 79–81
    - collapse in food web, 81–83
    - immature state, 83–84
    - lake fishery and restoration, 85
    - potential driving factors and underlying mechanisms, 84–85
- Ecosystem(s), 103–104
- functioning
    - changes, 79–81
    - evaluation, 74–76
  - model, 142–143
  - as object for modeling, 9–10
  - properties, 104–108
- Ecotox, 159
- Ecotoxicological models, 4, 6–7, 154, 203–205. *See also*
- Fugacity models
    - application, 141–144
    - application of models in, 154
    - characteristics, 144–153
      - dynamic model of toxic substance, 147–148
      - with effect components, 149–153
    - food chain, 146–147
    - in population dynamics, 148–149
    - static models of toxic substance mass flows, 147
    - contamination of agricultural products, 166–172
    - ecotoxicological properties, 159t
    - environmental properties of organic compounds, 158t
    - estimation of ecotoxicological parameters, 154–166
    - examples of toxic substance models, 155t–157t
      - as experimental tools, 172–176
      - heavy metal model, 172f
      - STELLA diagram of model, 175f
      - submodels of total ecotoxicological model, 151f
  - EFP. *See* Estimation of ecotoxicological parameters (EEP)
  - Effects models, 141–144
  - ELT. *See* Ecological law of thermodynamics (ELT)
  - Emergent wetlands, 234–236
  - EMF. *See* Ecological magnification factor (EMF)
  - Environmental
    - biogeochemical models, 33–37, 34f, 36t
    - factors, 136
    - management, 3, 3f
    - problems, 35, 37t
  - Environmental risk assessment (ERA), 142, 160
  - Equivalent indexes (EQIs), 92–93
  - ERA. *See* Environmental risk assessment (ERA)
  - ESLR. *See* Eustatic sea level rise (ESLR)
  - Estimation methods, 153
  - Estimation of ecotoxicological parameters (EEP), 165–166
    - network of estimation methods in, 166f
  - Euclidean motion, 92
  - Eustatic sea level rise (ESLR), 240–241
  - Experimental mathematics models, 171
  - Experimental tools, 4
    - ecotoxicological models as, 172–176
  - External variables, 16–18
- F**
- Fate models, 141–142
- Fate-transport-effect models (FTE models), 143, 203–205
- FCI. *See* Finn's cycling index (FCI)
- Fick's laws, 14–15
- Finn's cycling index (FCI), 74–76
- Finn's mean path length (FML), 74–76
- First law of population dynamics, 40–43
- Fla. *See* Fluoranthene (Fla)
- FLAM. *See* Fuzzy logic-based, adaptive management (FLAM)
- Flo. *See* Fluorene (Flo)
- Floating marsh, 238
- Fluoranthene (Fla), 187–188
- Fluorene (Flo), 187–188
- FML. *See* Finn's mean path length (FML)
- Food chain(s), 56–58, 56f, 146–147
  - bioaccumulation, 146f
  - model, 117, 118f
  - with two trophic levels and cycling of nutrients, 60–61

- Food web, 184  
collapse in, 81–83  
dynamic models, 146–147  
structure, 66, 77f
- Forcing functions, 16–18
- Forward propagation, 126–127
- FTE models. *See* Fate-transport-effect models (FTE models)
- FTESM. *See* Fundamental theorem of earth surface modeling (FTESM)
- FTS. *See* Fundamental theorem of surfaces (FTS)
- Fugacity models, 6–7, 182. *See also* Ecotoxicological models; LAVegMod vegetation model development and applications, 182–184  
future perspectives, 185  
for PAHs, 185–205
- Fundamental theorem of earth surface modeling (FTESM), 92  
and corollaries, 97–98
- Fundamental theorem of surfaces (FTS), 92
- Fuzzy adaptive management  
AM strategy, 224  
coupled natural and human systems, 212  
FLAM model, 222  
fuzzy TOPSIS, 223  
methods, 213–222  
AM, 215–216, 216f  
attributes, 214–215  
determining preferred management actions, 216–222  
estimating attributes, 216  
FLAM model for CNP, 213–214, 214f  
Linguistic rating scale, 218t  
triangular probability distribution, 218f
- Fuzzy decision rules, 212–213
- Fuzzy logic-based, adaptive management (FLAM), 213, 216. *See also* Fuzzy adaptive management model for CNP, 213–215, 214f
- Fuzzy models, 6–7
- Fuzzy Technique for Order Preference by Similarity of Ideal Solution (fuzzy TOPSIS), 213, 216–217, 219
- FVCOM, 245  
FVCOM-WASP model, 245
- G**
- Gaseous samples, 187–188
- GCMS. *See* Global climate models (GCMS)
- GCOMS. *See* Global Coastal Ocean Modelling System (GCOMS)
- Geographical information system (GIS), 91–92
- Geospiza fortis* (*G. fortis*), 115
- GIS. *See* Geographical information system (GIS)
- Global climate models (GCMS), 228–229
- Global Coastal Ocean Modelling System (GCOMS), 250
- Global models, 142–143
- Global warming, 104
- Gulf of Mexico Hypoxia, climate change impacts on, 243–245  
hypoxia models, 244–245  
model results, 245
- H**
- Hardwood forest, 236–237
- HASM. *See* High accuracy surface modeling (HASM)
- HCHs. *See* Hexachlorocyclohexanes (HCHs)
- Henry's constant, 161
- Henry's law, 189
- HERO chemicals. *See* Highly expected regulatory output chemicals (HERO chemicals)
- Hexachlorocyclohexanes (HCHs), 183
- High accuracy surface modeling (HASM), 92–97
- High-molecular-weight PAHs (HMW-PAHs), 195
- Highly expected regulatory output chemicals (HERO chemicals), 143
- HMW-PAHs. *See* High-molecular-weight PAHs (HMW-PAHs)
- Hold-out procedure, 128
- Hydrolysis, 173
- Hypoxia, 253–254  
models, 244–245
- I**
- IBM. *See* Individual-based models (IBM)
- IcdP. *See* Indeno[1,2,3-cd]pyrene (IcdP)
- ICM. *See* Integrated compartment model (ICM)
- IDW. *See* Inverse distance weighting (IDW)
- Indeno[1,2,3-cd]pyrene (IcdP), 187–188
- Individual-based models (IBM), 6–7
- Information eco-exergy, 111
- Integrated compartment model (ICM), 230
- Intergovernmental Panel on Climate Change (IPCC), 213  
Inverse distance weighting (IDW), 98–99
- IPCC. *See* Intergovernmental Panel on Climate Change (IPCC)
- J**
- Jackknife procedure, 129
- K**
- K-fold cross-validation, 128
- L**
- Lake Baiyangdian in Northern China  
fugacity models for PAHs, 185–205  
conceptual framework, 188

- ecological implications of proposed model, 203–205  
 equations for calculating fugacity capacities, 190t  
 equations for calculating transfer rate coefficients, 190t  
 model equations, 188  
 model parameters, 189  
 model validation, 196–200  
 modeled concentration distributions, 195–196  
 multimedia modeling, 193  
 parameter values for PAH components, 194t  
 QWASI fugacity model, 188f  
 sensitivities of modeled concentrations, 203  
 sensitivity analysis, 193–195  
 study area and measurements, 186–188  
 transfer fluxes, 200, 201t, 202f  
 uncertainty analysis, 195  
 uncertainty of modeled concentrations, 203
- Lake Small Baiyangdian and sampling sites, 187f
- Lake Chaohu, 70, 70f, 72t  
 diet composition, 75t  
 ecosystem properties, 80t  
 food web structure and relative biomasses, 77f  
 mixed trophic impacts, 82f  
 total fishery yield, 71f  
 trophic flows, 78f
- Lake fishery and restoration, 85
- Lake management, 69
- LAVegMod vegetation model, 230. *See also*  
 Ecotoxicological models; Fugacity models  
 barrier island vegetation, 239  
 development and implementation, 230–239, 231t, 232f, 232t  
 emergent wetlands, 234–236  
 floating marsh, 238  
 hardwood forest, 236–237  
 structure, 230–234  
 submerged aquatics, 238–239  
 swamp forest, 237–238
- LCMP. *See* Louisiana Coastal Master Plan (LCMP)
- Lead  
 balance of average Danish agriculture land, 167f  
 contamination of agricultural products by, 166–172
- Learning rate coefficient, 127–128
- Leave-one-out procedure, 129
- Limnoperna lacustris* (*L. lacustris*), 81–82
- Limnoithona sinensis* (*L. sinensis*), 81–82
- Linguistic rating scale, 218t
- LMW-PAHs. *See* Low-molecular-weight PAHs (LMW-PAHs)
- Logarithmic transformation, 130–131
- Logistic populations, 43, 44f  
 in periodic environment, 44–45, 46f  
 in random environment, 46–48
- Louisiana Coastal Master Plan (LCMP), 230
- Louisiana's coastal wetland plant communities, climate change effects on, 230–243  
 climate change scenarios, 240–241, 241t  
 LAVegMod model, 230–239, 231t, 232f, 232t  
 model inputs, 239–240, 240t  
 model results, 241–243, 242f–243f
- Low-molecular-weight PAHs (LMW-PAHs), 195
- ## M
- MAC. *See* Maximum allowable concentration (MAC)
- MacC. *See* Macrocrustacean crab (MacC)
- Macrobrachium nipponensis* (*M. nipponensis*), 81–82
- Macrocrustacean crab (MacC), 81–82
- Macrocrustacean shrimp (MacS), 73–74, 81–82
- Macrophytes, 205  
 samples, 187–188
- MacS. *See* Macrocrustacean shrimp (MacS)
- Malthus law, 40–43
- Mass conservation principles, 13–14
- Mathematical equations, 16–18
- Mathematical model, 40
- Maximum allowable concentration (MAC), 154–158
- Maximum loss index (MLI), 221
- Maximum sustainable average yield (MSAY), 45, 47
- Maximum sustainable yield (MSY), 43–44, 55
- McKay-type models, 142
- Metronidazole, 176
- Michaelis–Menten constant, 14–15
- Microcystis aeruginosa* (*M. aeruginosa*), 83
- Middle-molecular-weight PAHs (MMW-PAHs), 195
- Min–max normalization. *See* Range normalization
- Mixed trophic impacts analysis (MTI analysis), 74–76
- MLI. *See* Maximum loss index (MLI)
- MLP. *See* Multilayer perception (MLP)
- MMW-PAHs. *See* Middle-molecular-weight PAHs (MMW-PAHs)
- Modeled concentration distributions, 195–196
- Momentum coefficient, 127–128
- Monsoon dynamics, 247–248
- MSAY. *See* Maximum sustainable average yield (MSAY)
- MSY. *See* Maximum sustainable yield (MSY)
- MTI analysis. *See* Mixed trophic impacts analysis (MTI analysis)
- Multilayer perception (MLP), 125  
 advantages, 133  
 BP learning algorithm, 126–128  
 data preprocessing, 130–131  
 disadvantages, 133  
 in ecological modeling, 132  
 overfitting, 131  
 recommendations for ecological modeling, 134  
 sensitivity analysis, 131–132  
 structure, 125–126  
 three-layered feed-forward neural network, 125f

- Multilayer perception (MLP) (*Continued*)  
 usage in ecological modeling, 134–138  
 validation of models, 128–129
- Multimedia modeling, 193
- Mutualism, 55, 56f
- N**
- NASA Earth Exchange (NEX), 222–223
- NEC. *See* Noneffect concentrations (NEC)
- Net system production (NSP), 74–76
- Neurons, 125–126
- NEX. *See* NASA Earth Exchange (NEX)
- NGOM. *See* Northern Gulf of Mexico shelf (NGOM)
- NOEC. *See* Nonobserved-effect concentration (NOEC)
- Noneffect concentrations (NEC), 154–158
- Nonlinear computational elements, 125–126
- Nonobserved-effect concentration (NOEC), 142
- Northern Gulf of Mexico shelf (NGOM), 243–244  
 hypoxia in, 244
- NPZD. *See* Nutrient-Phytoplankton-Zooplankton-Detritus (NPZD)
- NSP. *See* Net system production (NSP)
- Numerical models, 228–229
- Numerical simulation models, 228
- Nutrient cycling, 59–60, 59f  
 food chain with, 60–61
- Nutrient-Phytoplankton-Zooplankton-Detritus (NPZD), 251
- Nyssa aquatica, 237
- O**
- OCPs. *See* Organochlorine pesticides (OCPs)
- Ordinary Kriging (OK), 98–99
- Organic compounds, environmental properties of, 158t
- Organism models, 142–143
- Organochlorine pesticides (OCPs), 183
- Overfitting, 131
- P**
- P/B ratio. *See* Production/biomass ratio (P/B ratio)
- PAHs. *See* Polycyclic aromatic hydrocarbon (PAHs)
- Palaemon modestus* (*P. modestus*), 76
- Parameter  
 ecological model, 16–18  
 estimation methods, 28–29
- Partial derivatives method, 131–132
- Particulate samples, 187–188
- PCBs. *See* Polychlorinated biphenyls (PCBs)
- PEC. *See* Predicted environmental concentration (PEC)
- Periodic environment, logistic population in, 44–45, 46f
- Perturbation method, 131–132
- Phenanthrene (Phe), 186
- Photolysis, 173
- Phytoplankton, 27–28, 59–60
- Political/cultural model, 153
- Polychlorinated biphenyls (PCBs), 183–184
- Polycyclic aromatic hydrocarbon (PAHs), 183, 186  
 in bulk and subphases, 196t  
 comparisons between subphases, 199f  
 fugacity model for, 185–205  
 conceptual framework, 188  
 ecological implications of proposed model, 203–205  
 equations for calculating fugacity capacities, 190t  
 equations for calculating transfer rate coefficients, 190t  
 model equations, 188  
 model parameters, 189  
 model validation, 196–200  
 modeled concentration distributions, 195–196  
 multimedia modeling, 193  
 parameter values for PAH components, 194t  
 QWASI fugacity model, 188f  
 sensitivities of modeled concentrations, 203  
 sensitivity analysis, 193–195  
 study area and measurements, 186–188  
 transfer fluxes, 200, 201t, 202f  
 uncertainty analysis, 195  
 uncertainty of modeled concentrations, 203  
 in subphases in Lake Small Baiyangdian, 198f
- Polymer compounds, 163–164
- Population conservation equations, 40
- Population dynamics  
 ecotoxicological models in, 148–149  
 models, 6–7, 117, 118f
- Population models, 39–40, 142–143
- Potential driving factors and underlying mechanisms, 84–85
- Predicted environmental concentration (PEC), 142
- Predictive model, 40
- Preferred management actions, 216–222  
 best adaptive management strategy, 222  
 for climate change scenarios, 217–220  
 for time periods, 221–222
- Prey–predator models, 48–52, 48f, 50f  
 Rozenzweig–McArthur model, 51f
- Primary biodegradation, 163
- Process rate equations, 15t
- Process-functional approach, 9
- Processing units, 125–126
- Production/biomass ratio (P/B ratio), 73–74
- Profile method, 131–132
- Pyrene (Pyr), 187–188
- Q**
- Q/B ratio. *See* Consumption/biomass ratio (Q/B ratio)
- Quantitative watereair sediment interaction (QWASI), 186

- fugacity model, 188, 188f  
 mass balance equations in, 189t  
 parameter values for PAH components in, 194t  
 parameters for, 191t–193t  
 sensitivity coefficients of sensitive parameters, 204t  
 transfer and transformation processes, 189t
- R**
- Random environment, logistic population in, 46–48  
 dynamics of two simulations, 47f
- Range normalization, 130–131
- Rational actor model, 153
- RCPs. *See* Representative concentration pathways (RCPs)
- Regional Ocean Modeling System (ROMS), 248, 251
- Remote sensing (RS), 91–92
- Representative concentration pathways (RCPs), 213
- Risk management systems, 153
- ROMS. *See* Regional Ocean Modeling System (ROMS)
- RS. *See* Remote sensing (RS)
- S**
- Salix nigra, 237
- SAV. *See* Submerged aquatic vegetation (SAV)
- Scientific hypotheses, 4
- SD. *See* Standard deviation (SD)
- SDMs. *See* Structurally dynamic models (SDMs)
- SE rates. *See* Soil erosion rates (SE rates)
- Second law of population dynamics, 43–44
- Sediment transport, 248–250
- Self-organizing maps (SOM), 6–7, 125
- Semisulcospira cancellata* (*S. cancellata*), 81–82
- Sensitivity analysis, 24–27, 27t, 131–132, 193–195
- Shifts in species composition, 106, 106f, 117
- Simulate WAve Nearshore model (SWAN), 248
- Sinocalanus dorii* (*S. dorii*), 81–82
- SOI. *See* System omnivory index (SOI)
- Soil erosion rates (SE rates), 215
- Soil morphology model, 240
- SOM. *See* Self-organizing maps (SOM)
- Spatial distribution, 5
- Spatial models, 6–7
- Species, 106–107
- SSC. *See* Suspended sediment concentration (SSC)
- SSEs. *See* Sum of square errors (SSEs)
- Standard deviation (SD), 176
- Standardization, 130–131
- State variables, 16–18
- Static models of toxic substance mass flows, 147
- Statistical analysis, 1–2
- Steady state models, 6–7, 66. *See also* Biogeochemical models; Dynamic population models  
 ecopath model, 66–67  
 future perspectives, 68
- STELLA software, 24, 168  
 Cd model, 169f  
 heavy metal model, 172f  
 STELLA diagram of model, 175f
- Stochastic decision rules, 212
- Stochastic models, 6–7
- Structurally dynamic models (SDMs), 6–7, 103–104, 107  
 application for assessment of ecological changes, 117–121  
 biomass and work energy, 119f–120f  
 biomass for population, 119t  
 food chain model, 118f  
 population dynamic model, 118f  
 development, 108–112, 110f  
 for Darwin's finches, 115–116  
 ecosystem properties, 104–108  
 using WE as goal function, 112–115
- Submerged aquatic vegetation (SAV), 233, 238–239
- Sulfamethoxazole, 176
- Sum of square errors (SSEs), 135, 135f
- Supervised learning, 125
- Surface modeling, 91–92
- Suspended sediment concentration (SSC), 249f
- Swamp forest, 237–238
- SWAN. *See* Simulate WAve Nearshore model (SWAN)
- System omnivory index (SOI), 74–76
- T**
- Tapertail anchovy (TapA), 81
- TC. *See* Total consumption (TC)
- Tcy. *See* Throughput cycled (Tcy)
- TD. *See* Total detritus (TD)
- Tentative modeling procedure, 19–20, 21f
- Terrestrial vegetation, 184
- TEEx. *See* Total exports (TEEx)
- Thermodynamic equilibrium, 109–111
- Thermodynamic translation of Darwin's theory, 107–108
- Throughput cycled (Tcy), 74–76
- Time periods, preferred management actions for, 221–222
- TIN. *See* Triangulated irregular network (TIN)
- TLs. *See* Trophic levels (TLs)
- Total consumption (TC), 74–76
- Total detritus (TD), 74–76
- Total exports (TEEx), 74–76
- Total net primary production (TPP), 74–76
- Total net primary production and total biomass (TPP/TB), 74–76
- Total net primary production with total respiration (TPP/TR), 74–76
- Total respiratory flows (TR flows), 74–76
- Total system throughput (TST), 74–76

Toxic substance  
  dynamic model, 147–148  
  models, 144–145, 155t–157t  
  at one trophic level, 148f  
  in organism *vs.* time, 150f  
  static models of toxic substance mass flows, 147  
TPP. *See* Total net primary production (TPP)  
TPP/TB. *See* Total net primary production and total biomass (TPP/TB)  
TPP/TR. *See* Total net primary production with total respiration (TPP/TR)  
TR flows. *See* Total respiratory flows (TR flows)  
Transfer fluxes of PAHs, 200, 201t, 202f  
Trial-and-error calibration, 29–30  
Triangular probability distribution, 218f  
Triangulated irregular network (TIN), 98–99  
Trophic levels (TLs), 69  
  determination, 76  
  different functional groups in Lake Chaohu, 79f  
TST. *See* Total system throughput (TST)  
Two-tailed *t*-test, 176

## U

Uncertainty analysis, 195  
Unsupervised learning, 125

## V

Validation, 19, 32  
  of models, 128–129

Variable parameter models, 143  
Variance normalization. *See* Standardization  
Vegetation, 184  
Verhulst law, 43–44  
Verification, 19, 24

## W

Wastewater stabilization pond system (WSP), 173  
  removal of antibiotic from, 174f  
Water solubility, 161  
Weather Research Forecast model (WRF), 248  
Weights method, 131–132  
Western Pacific, climate change impacts on coastal ecosystems in, 245–250  
  coastal ecosystem processes, 250  
  monsoon dynamics, 247–248  
  sediment transport, 248–250  
Work energy (WE), 107–109, 109f, 117  
  as goal function in SDM, 112–115

## X

Xenobiotic compounds, microbiological decomposition of, 163

## Y

Yellowstone National Park, 212

## Z

Zooplankton, 27–28, 81–82