

# Universo Primitivo





## 2022-2023 (1º Semestre)

Mestrado em Física - Astronomia



### Chapter 2

#### 2. The Standard Model of Cosmology (SMC)

- Fundamental assumptions;
- The GR equations and the Friedmann-Lemaitre-Robertson-Walker (FLRW) solution;
- FLRW models:
  - Dynamic equations;
  - Energy-momentum conservation;
  - Fluid components and equations of state;
  - Cosmological parameters;
  - The Friedmann equation: the evolutionary phases of the Universe; exact solutions: age of the Universe;
  - Distances; horizons and volumes;
  - The accelerated expansion of the Universe;
- Problems with the SMC: Horizon; Flatness; Relic particles; origin of perturbations; primordial Isotropy and homogeneity
- The idea of Inflation



Ch. 1



Ch. 1

3

# Standard Model of Cosmology

# SMC: Mathematical framework

Fundamental assumptions:

- The Universe is homogeneous and isotropic when observed on large scales and expands uniformly with respect to any position
- The dynamics of space-time is described by Einstein's theory of general relativity (GR).

$$G_{ab} = R_{ab} - \frac{1}{2}Rg_{ab} = \frac{8\pi G}{c^4}T_{ab}$$

for the Universe to be homogeneous and isotropic the stress-energy tensor must be that of a perfect fluid

$$T_{ab} = (\rho + \frac{p}{c^2})U_aU_b - \frac{p}{c^2}g_{ab}$$

5

# SMC: Mathematical framework

The cosmological constant in the GR equation:

$$G_{\mu\nu} + \Lambda g_{\mu\nu} = 8\pi G T_{\mu\nu}, \quad (\Lambda \text{ as "cosmological constant"})$$

$$G_{\mu\nu} = 8\pi G \left( T_{\mu\nu} - \frac{\Lambda}{8\pi G} g_{\mu\nu} \right) = 8\pi G \tilde{T}_{\mu\nu}, \quad (\Lambda \text{ as "vacuum energy"})$$

The Einstein tensor, Ricci tensor and Ricci scalar are:

$$G_{\mu\nu} = R_{\mu\nu} - \frac{1}{2}g_{\mu\nu}R$$

$$R_{\mu\nu} = \Gamma_{\mu\nu,\alpha}^{\alpha} - \Gamma_{\mu\alpha,\nu}^{\alpha} + \Gamma_{\mu\nu}^{\alpha}\Gamma_{\alpha\beta}^{\beta} - \Gamma_{\mu\beta}^{\alpha}\Gamma_{\alpha\nu}^{\beta}$$

$$R = g^{\mu\nu}R_{\mu\nu}$$

$$\Gamma_{\nu\lambda}^{\mu} = \frac{1}{2}g^{\mu\nu}(g_{\alpha\nu,\lambda} + g_{\alpha\lambda,\nu} - g_{\nu\lambda,\alpha}) \quad g_{\mu\nu,\lambda} \equiv \partial g_{\alpha\nu}/\partial x^{\lambda} \quad g^{\mu\lambda}g_{\lambda\nu} = \delta_{\nu}^{\mu}$$

where,


$$ds^2 = \sum_{\mu,\nu=0}^3 g_{\mu\nu}dX^{\mu}dX^{\nu} \equiv g_{\mu\nu}dX^{\mu}dX^{\nu}$$

6


# SMC: Mathematical framework

## Einstein Equation:


$$G_{\mu\nu} + \Lambda g_{\mu\nu} = 8\pi G T_{\mu\nu},$$



Albert Einstein  
1879-1955



More accurate ways of visualizing GR: <https://youtu.be/wrwgIjBUYVc>



# SMC: Mathematical framework

## Geodesic Equation:

In the absence of non-gravitational forces, free falling particles move along “geodesics”, described by the so-called Geodesic equation.

$$\frac{dU^\mu}{ds} + \Gamma_{\alpha\beta}^\mu U^\alpha U^\beta = 0$$

where,

$$U^\mu \equiv \frac{dX^\mu}{ds}$$

four-velocity of the particle along its free-falling path  $X^\mu(s)$

$$S = -m \int_A^B ds .$$




Figure 1.4: Parameterisation of an arbitrary path in spacetime,  $X^\mu(\lambda)$ .

# SMC: Mathematical framework

Fundamental assumptions:

- The Universe is homogeneous and isotropic when observed on large scales and expands uniformly with respect to any position
- The dynamics of space-time is described by Einstein's theory of general relativity (GR).

$$G_{ab} = R_{ab} - \frac{1}{2}Rg_{ab} = \frac{8\pi G}{c^4}T_{ab} - \Lambda g_{ab} \quad T_{ab} = \left(\rho + \frac{p}{c^2}\right)U_aU_b - \frac{p}{c^2}g_{ab}$$

In these conditions **the solution of the Einstein equation** is the Friedmann-Lemaitre-Robertson-Walker (**FLRW**) metric:

$$ds^2 = c^2dt^2 - a^2(t) \left[ \frac{dr^2}{1 - kr^2} + r^2 (d\theta^2 + \sin^2\theta d\phi^2) \right] \quad 9$$

# SMC: Mathematical framework

- Dynamical equations:  
(result from the Einstein equations and govern the time evolution of  $a(t)$ )

$$\left(\frac{\dot{a}}{a}\right)^2 = \frac{8\pi G}{3}\rho + \frac{\Lambda c^2}{3} - \frac{kc^2}{a^2}$$

$$\frac{\ddot{a}}{a} = -\frac{4\pi G}{3}\left(\rho + 3\frac{p}{c^2}\right) + \frac{\Lambda c^2}{3}$$

Friedmann equation

Raychaudhuri  
(or acceleration) equation

- Energy momentum conservation:  $\nabla_\mu T^\mu_\nu \equiv T^\mu_{\nu;\mu} = 0$

the covariant derivative reads:  $\nabla_\mu T^\mu_\nu = \partial_\mu T^\mu_\nu + \Gamma^\mu_{\mu\lambda} T^\lambda_\nu - \Gamma^\lambda_{\mu\nu} T^\mu_\lambda = 0$

the  $\nu = 0$  (time) component of this equation gives:

$$\dot{\rho} = -3\frac{\dot{a}}{a}\left(\rho + \frac{p}{c^2}\right) \Rightarrow d(\rho c^2 a^3) = -pd(a^3) \quad \text{Energy conservation equation}$$

$$p = w\rho c^2 \quad -1 \leq w \leq 1 \quad \text{Equation of State (EoS)}$$

for fluids with constant EoS parameter,  $w$ , the solution is:

$$\rho(t) = \rho_i \left(\frac{a(t)}{a_i}\right)^{-3(1+w)}$$

# SMC: Mathematical framework

## Covariant derivative:

*Covariant derivative.*—The covariant derivative is an important object in differential geometry and it is of fundamental importance in general relativity. The geometrical meaning of  $\nabla_\mu$  will be discussed in detail in the GR course. In this course, we will have to be satisfied with treating it as an operator that acts in a specific way on scalars, vectors and tensors:

- There is no difference between the covariant derivative and the partial derivative if it acts on a scalar

$$\nabla_\mu f = \partial_\mu f . \quad (1.3.83)$$

- Acting on a contravariant vector,  $V^\nu$ , the covariant derivative is a partial derivative plus a correction that is linear in the vector:

$$\nabla_\mu V^\nu = \partial_\mu V^\nu + \Gamma^\nu_{\mu\lambda} V^\lambda . \quad (1.3.84)$$

Look carefully at the index structure of the second term. A similar definition applies to the covariant derivative of covariant vectors,  $\omega_\nu$ ,

$$\nabla_\mu \omega_\nu = \partial_\mu \omega_\nu - \Gamma^\lambda_{\mu\nu} \omega_\lambda . \quad (1.3.85)$$

Notice the change of the sign of the second term and the placement of the dummy index.

- For tensors with many indices, you just repeat (1.3.84) and (1.3.85) for each index. For each upper index you introduce a term with a single  $+\Gamma$ , and for each lower index a term with a single  $-\Gamma$ :

$$\begin{aligned} \nabla_\sigma T^{\mu_1 \mu_2 \dots \mu_k}_{\nu_1 \nu_2 \dots \nu_l} &= \partial_\sigma T^{\mu_1 \mu_2 \dots \mu_k}_{\nu_1 \nu_2 \dots \nu_l} \\ &+ \Gamma^{\mu_1}_{\sigma\lambda} T^{\lambda \mu_2 \dots \mu_k}_{\nu_1 \nu_2 \dots \nu_l} + \Gamma^{\mu_2}_{\sigma\lambda} T^{\mu_1 \lambda \dots \mu_k}_{\nu_1 \nu_2 \dots \nu_l} + \dots \\ &- \Gamma^\lambda_{\sigma\nu_1} T^{\mu_1 \mu_2 \dots \mu_k}_{\lambda \nu_2 \dots \nu_l} - \Gamma^\lambda_{\sigma\nu_2} T^{\mu_1 \mu_2 \dots \mu_k}_{\nu_1 \lambda \dots \nu_l} - \dots \end{aligned} \quad (1.3.86)$$

This is the general expression for the covariant derivative. Luckily, we will only be dealing with relatively simple tensors, so this monstrous expression will usually reduce to something manageable.

11

# SMC: Mathematical framework

- EoS for different energy density components:

- $w = 1/3$  (radiation)

$$\rho_\gamma = \rho_{\gamma 0} \left(\frac{a_0}{a}\right)^4 \xrightarrow{(1)} \left(\frac{\dot{a}}{a}\right)^2 \propto \frac{1}{a^4} \rightarrow a \propto t^{1/2}$$

$$\rho(t) = \rho_i \left(\frac{a(t)}{a_i}\right)^{-3(1+w)}$$

$$\left(\frac{\dot{a}}{a}\right)^2 = \frac{8\pi G}{3} \rho$$

- $w = 0$  (collisionless matter)

$$\rho_m = \rho_{m0} \left(\frac{a_0}{a}\right)^3 \xrightarrow{(2)} \left(\frac{\dot{a}}{a}\right)^2 \propto \frac{1}{a^3} \rightarrow a \propto t^{2/3}$$

- $w = -1$  (cosmological constant)

$$\rho_\Lambda = \Lambda/8\pi G = -P_\Lambda \xrightarrow{(3)} a \propto e^{\sqrt{\Lambda}t}$$

(1) after integration of the Friedmann equation with  $k = 0$ ,  $\Lambda = 0$ ,  $\rho = \rho_\gamma$ .

(2) after integration of the Friedmann equation with  $k = 0$ ,  $\Lambda = 0$ ,  $\rho = \rho_m$ .

(3) after integration of the Friedmann equation with  $k = 0$ ,  $\Lambda = 8\pi G\rho_\Lambda$ ,  $\rho = 0$

12

# SMC: FLRW models

- Cosmological parameters:

The Friedmann equation

$$\left(\frac{\dot{a}}{a}\right)^2 = \frac{8\pi G}{3}\rho + \frac{\Lambda c^2}{3} - \frac{kc^2}{a^2}$$

can be re-written as

$$H^2 = \frac{8\pi G}{3}(\rho_r + \rho_B + \rho_{DM}) + \frac{\Lambda c^2}{3} - \frac{kc^2}{a^2}$$

where,

$$H(t) = \frac{\dot{a}(t)}{a(t)}$$

$$\rho \equiv \sum_i \rho_i$$

dividing by  $H^2$  on gets

$$1 = \frac{8\pi G}{3H^2}\rho_r + \frac{8\pi G}{3H^2}\rho_B + \frac{8\pi G}{3H^2}\rho_{DM} + \frac{\Lambda c^2}{H^2} - \frac{kc^2}{a^2 H^2}$$

or

$$1 = \Omega_r + \Omega_B + \Omega_{DM} + \Omega_\Lambda + \Omega_k$$

13

# SMC: FLRW models

$$\left(\frac{\dot{a}}{a}\right)^2 = \frac{8\pi G}{3}\rho + \frac{\Lambda c^2}{3} - \frac{kc^2}{a^2}$$

- Cosmological parameters:

$$\frac{8\pi G}{3H^2}\rho + \frac{\Lambda c^2}{3H^2} - \frac{kc^2}{a^2 H^2} = 1 \Leftrightarrow \sum_i \Omega_i + \Omega_\Lambda + \Omega_k = 1$$

$$H(t) = \frac{\dot{a}(t)}{a(t)}$$

Hubble parameter

$$\Omega_i \equiv \frac{\rho_i}{\rho_{\text{crit}}}$$

Mater-energy density parameters:  
 $(\Omega = \frac{\rho}{\rho_c} = \sum \frac{\rho_i}{\rho_c} = \sum \Omega_i)$

$$\Omega_\Lambda = \frac{\Lambda c^2}{3H^2}$$


Vacuum or dark energy density parameter

$$\Omega_k = -\frac{kc^2}{a^2 H^2}$$

Curvature density parameter

$$\rho_c = \frac{3H^2}{8\pi G}$$

Critical energy density



$$\rho \equiv \sum_i \rho_i$$

$$\Omega_i \equiv \frac{\rho_i}{\rho_{\text{crit}}}$$

includes all matter and radiation components (baryons, dark mater, radiation, ...)


# SMC: FLRW models

- Friedmann equation revisited

$$H^2(t) = \frac{8\pi G}{3}(\rho_r + \rho_m) - \frac{kc^2}{a^2} + \frac{\Lambda c^2}{3}$$

$$= H_0^2 \left[ \Omega_{r0} \left(\frac{a_0}{a}\right)^4 + \Omega_{m0} \left(\frac{a_0}{a}\right)^3 + \Omega_{k0} \left(\frac{a_0}{a}\right)^2 + \Omega_{\Lambda 0} \right]$$

The evolutionary fate of the Universe is determined by cosmological parameters



## SMC: Exact solutions of the Friedmann equation

- Scale factor:

$$\left(\frac{\dot{a}}{a}\right)^2 = \frac{8\pi G}{3}\rho + \frac{\Lambda c^2}{3} - \frac{kc^2}{a^2}$$

$$\frac{d a(t)}{dt a_0} = H_0 \sqrt{1 - \Omega_0 + \Omega_{m0} \left(\frac{a}{a_0}\right)^{-1} + \Omega_{r0} \left(\frac{a}{a_0}\right)^{-2} - \Omega_{\Lambda 0} \left[1 - \left(\frac{a}{a_0}\right)^2\right]}$$

for a critical density ( $\Omega_k = \Omega_\Lambda = 0$ ) universe, gives:

$$\frac{a(t)}{a_0} = \left(\frac{3(1+w)}{2} H_0 t\right)^{2/(3(1+w))}$$

$$H(t) = \frac{\dot{a}}{a} = \frac{2}{3(w+1)t}$$

- Age of the Universe:

$$t = H_0^{-1} \int_0^{\frac{a(t)}{a_0} = (1+z)^{-1}} \frac{1}{\sqrt{1 - \Omega_0 + \Omega_{m0}x^{-1} + \Omega_{r0}x^{-2} - \Omega_\Lambda (1 - x^2)}} dx$$

Redshift:

$$z = \frac{E - E_0}{E_0} =$$

$$= \frac{v}{v_0} - 1 =$$


$$= \frac{\lambda_0}{\lambda} - 1 =$$

$$= \frac{a_0}{a} - 1$$



# SMC: Concordance Cosmology


Combination of different observational datasets...



From:

... allow us to impose constraints on cosmological parameters

$$\sum_i \Omega_i + \Omega_\Lambda + \Omega_k = 1$$



## WMAP3 parameters

Parameter	Value	Description
<i>Basic parameters</i>		
$H_0$	$70.9^{+2.4}_{-3.2} \text{ km s}^{-1} \text{ Mpc}^{-1}$	Hubble parameter
$\Omega_b$	$0.0444^{+0.0042}_{-0.0035}$	Baryon density
$\Omega_m$	$0.266^{+0.025}_{-0.040}$	Total matter density (baryons + dark matter)
$\tau$	$0.079^{+0.029}_{-0.032}$	Optical depth to reionization
$A_s$	$0.813^{+0.042}_{-0.052}$	Scalar fluctuation amplitude
$n_s$	$0.948^{+0.015}_{-0.018}$	Scalar spectral index
<i>Derived parameters</i>		
$\rho_0$	$0.94^{+0.06}_{-0.09} \times 10^{-26} \text{ kg/m}^3$	Critical density
$\Omega_\Lambda$	$0.732^{+0.040}_{-0.025}$	Dark energy density
$z_{\text{ion}}$	$10.5^{+2.6}_{-2.9}$	Reionization red-shift
$\sigma_8$	$0.772^{+0.036}_{-0.048}$	Galaxy fluctuation amplitude
$t_0$	$13.73^{+0.13}_{-0.17} \times 10^9 \text{ years}$	Age of the universe



# SMC: Cosmological parameters after Planck

From: Planck collaboration. XVI. arXiv:1303.5076

**Table 2.** Cosmological parameter values for the six-parameter base  $\Lambda$ CDM model. Columns 2 and 3 give results for the *Planck* temperature power spectrum data alone. Columns 4 and 5 combine the *Planck* temperature data with *Planck* lensing, and columns 6 and 7 include *WMAP* polarization at low multipoles. We give best fit parameters (i.e. the parameters that maximise the overall likelihood for each data combination) as well as 68% confidence limits for constrained parameters. The first six parameters have flat priors. The remainder are derived parameters as discussed in Sect. 2. Beam, calibration parameters, and foreground parameters (see Sect. 4) are not listed for brevity. Constraints on foreground parameters for *Planck*+WP are given later in Table 5.


Parameter	<i>Planck</i>		<i>Planck</i> +lensing		<i>Planck</i> +WP	
	Best fit	68% limits	Best fit	68% limits	Best fit	68% limits
$\Omega_b h^2$	0.022068	$0.02207 \pm 0.00033$	0.022242	$0.02217 \pm 0.00033$	0.022032	$0.02205 \pm 0.00028$
$\Omega_c h^2$	0.12029	$0.1196 \pm 0.0031$	0.11805	$0.1186 \pm 0.0031$	0.12038	$0.1199 \pm 0.0027$
$100\theta_{\text{MC}}$	1.04122	$1.04132 \pm 0.00068$	1.04150	$1.04141 \pm 0.00067$	1.04119	$1.04131 \pm 0.00063$
$\tau$	0.0925	$0.097 \pm 0.038$	0.0949	$0.089 \pm 0.032$	0.0925	$0.089^{+0.012}_{-0.014}$
$n_s$	0.9624	$0.9616 \pm 0.0094$	0.9675	$0.9635 \pm 0.0094$	0.9619	$0.9603 \pm 0.0073$
$\ln(10^{10} A_s)$	3.098	$3.103 \pm 0.072$	3.098	$3.085 \pm 0.057$	3.0980	$3.089^{+0.024}_{-0.027}$
$\Omega_\Lambda$	0.6825	$0.686 \pm 0.020$	0.6964	$0.693 \pm 0.019$	0.6817	$0.685^{+0.018}_{-0.016}$
$\Omega_m$	0.3175	$0.314 \pm 0.020$	0.3036	$0.307 \pm 0.019$	0.3183	$0.315^{+0.016}_{-0.018}$
$\sigma_8$	0.8344	$0.834 \pm 0.027$	0.8285	$0.823 \pm 0.018$	0.8347	$0.829 \pm 0.012$
$z_{\text{re}}$	11.35	$11.4^{+4.0}_{-2.8}$	11.45	$10.8^{+3.1}_{-2.5}$	11.37	$11.1 \pm 1.1$
$H_0$	67.11	$67.4 \pm 1.4$	68.14	$67.9 \pm 1.5$	67.04	$67.3 \pm 1.2$
$10^9 A_s$	2.215	$2.23 \pm 0.16$	2.215	$2.19^{+0.12}_{-0.14}$	2.215	$2.196^{+0.051}_{-0.060}$
$\Omega_m h^2$	0.14300	$0.1423 \pm 0.0029$	0.14094	$0.1414 \pm 0.0029$	0.14305	$0.1426 \pm 0.0025$
$\Omega_b h^3$	0.09597	$0.09590 \pm 0.00059$	0.09603	$0.09593 \pm 0.00058$	0.09591	$0.09589 \pm 0.00057$
$Y_p$	0.247710	$0.24771 \pm 0.00014$	0.247785	$0.24775 \pm 0.00014$	0.247695	$0.24770 \pm 0.00012$
Age/Gyr	13.819	$13.813 \pm 0.058$	13.784	$13.796 \pm 0.058$	13.8242	$13.817 \pm 0.048$
$z_*$	1090.43	$1090.37 \pm 0.65$	1090.01	$1090.16 \pm 0.65$	1090.48	$1090.43 \pm 0.54$
$r_*$	144.58	$144.75 \pm 0.66$	145.02	$144.96 \pm 0.66$	144.58	$144.71 \pm 0.60$
$100\theta_*$	1.04139	$1.04148 \pm 0.00066$	1.04164	$1.04156 \pm 0.00066$	1.04136	$1.04147 \pm 0.00062$
$z_{\text{drag}}$	1059.32	$1059.29 \pm 0.65$	1059.59	$1059.43 \pm 0.64$	1059.25	$1059.25 \pm 0.58$
$r_{\text{drag}}$	147.34	$147.53 \pm 0.64$	147.74	$147.70 \pm 0.63$	147.36	$147.49 \pm 0.59$
$k_{\text{D}}$	0.14026	$0.14007 \pm 0.00064$	0.13998	$0.13996 \pm 0.00062$	0.14022	$0.14009 \pm 0.00063$
$100\theta_{\text{D}}$	0.161332	$0.16137 \pm 0.00037$	0.161196	$0.16129 \pm 0.00036$	0.161375	$0.16140 \pm 0.00034$
$z_{\text{eq}}$	3402	$3386 \pm 69$	3352	$3362 \pm 69$	3403	$3391 \pm 60$
$100\theta_{\text{eq}}$	0.8128	$0.816 \pm 0.013$	0.8224	$0.821 \pm 0.013$	0.8125	$0.815 \pm 0.011$
$r_{\text{drag}}/D_V(0.57)$	0.07130	$0.0716 \pm 0.0011$	0.07207	$0.0719 \pm 0.0011$	0.07126	$0.07147 \pm 0.00091$

# SMC: Cosmological parameters after Planck



# SMC: Cosmological parameters after Planck

$$1 = \Omega_r + \Omega_B + \Omega_{DM} + \Omega_\Lambda + \Omega_k$$





Before Planck

After Planck

# SMC: Cosmological parameters after Planck


From: Planck collaboration. XVI. arXiv:1303.5076






## SMC: Limitations of a 6-parameter model...



Comparing primary CMB with other datasets




- Higher values of  $\Omega_m, \sigma_8$  in *Planck* CMB analysis
- $3\sigma$  tension
- More general tension between clusters and CMB ?





# SMC: Limitations of a 6 parameter model...



## Comparing primary CMB with other datasets



- Getting higher  $\sigma_8$  from clusters
- Change scaling
- Change bias
- Account for missing clusters
- Getting lower  $\sigma_8$  from CMB
- Change initial power spectrum
- Change transfer function



M. Douspis, 03/04/2013, Cosmology from Planck SZ cluster counts

14


## Planck Legacy: A new baseline cosmological model?

### The (new) concordance model: $\Lambda$ CDM + massive neutrinos

From: Planck collaboration. XIII (2015)

Parameter	TT	TT+lensing	TT+lensing+ext	TT, TE, EE	TT, TE, EE+lensing	TT, TE, EE+lensing+ext
$\Omega_b$	$-0.052^{+0.049}_{-0.066}$	$-0.005^{+0.016}_{-0.012}$	$-0.0001^{+0.0054}_{-0.0052}$	$-0.040^{+0.038}_{-0.041}$	$-0.004^{+0.015}_{-0.012}$	$0.0008^{+0.0040}_{-0.0029}$
$\Sigma m_\nu$ [eV]	$< 0.715$	$< 0.675$	$< 0.234$	$< 0.492$	$< 0.589$	$< 0.194$
$N_{\text{eff}}$	$3.12^{+0.64}_{-0.63}$	$3.12^{+0.62}_{-0.61}$	$3.15^{+0.41}_{-0.40}$	$2.99^{+0.41}_{-0.39}$	$2.94^{+0.38}_{-0.38}$	$3.04^{+0.33}_{-0.33}$
$Y_p$	$0.252^{+0.041}_{-0.042}$	$0.251^{+0.040}_{-0.039}$	$0.251^{+0.035}_{-0.036}$	$0.250^{+0.026}_{-0.027}$	$0.247^{+0.026}_{-0.027}$	$0.249^{+0.025}_{-0.026}$
$dn_s/d \ln k$	$-0.008^{+0.016}_{-0.016}$	$-0.003^{+0.015}_{-0.015}$	$-0.003^{+0.015}_{-0.014}$	$-0.006^{+0.014}_{-0.014}$	$-0.002^{+0.013}_{-0.013}$	$-0.002^{+0.013}_{-0.013}$
$r_{0.002}$	$< 0.103$	$< 0.114$	$< 0.114$	$< 0.0987$	$< 0.112$	$< 0.113$
$w$	$-1.54^{+0.62}_{-0.50}$	$-1.41^{+0.64}_{-0.56}$	$-1.006^{+0.085}_{-0.091}$	$-1.55^{+0.58}_{-0.48}$	$-1.42^{+0.62}_{-0.56}$	$-1.019^{+0.075}_{-0.080}$

$$\sum m_\nu = 0.16^{+0.08}_{-0.11} \text{ eV} \quad (\text{Planck TT+lowP+aggressive lensing + BAO; 68\%})$$



# SMC: Particle and Event horizons

Consider light travelling along radial ( $d\theta = d\phi = 0$ ) geodesics in a FLRW metric ( $c=1$ ):

$$\begin{aligned}
 ds^2 &= dt^2 - a^2(t) \left[ \frac{dr^2}{1 - kr^2} + r^2(d\theta^2 + \sin^2\theta d\phi^2) \right], \\
 &= dt^2 - a^2(t) [d\chi^2 + f_k(\chi)(d\theta^2 + \sin^2\theta d\phi^2)],
 \end{aligned}$$

( $d\chi = dr$  for flat geometries, see eg Sec. 1.2.3 Baumann). Let's set  $d\theta = d\phi = 0$  and define **conformal time** as  $d\tau = dt/a$ . This allows us to write:

$$ds^2 = a^2(\tau) [d\tau^2 - d\chi^2]$$



Since **light rays travel along null** ( $ds^2 = 0$ ) **geodesics**:  $d\chi = \pm d\tau$

Integrating from the **past** ( $t_i$ ) to **present** ( $t$ ) or from **the present to the future** ( $t_f$ ) one can define:

- **Particle horizon**:  $\chi_{ph}(\tau) = \tau - \tau_i = \int_{t_i}^t \frac{dt}{a(t)}$  with  $t_i = 0$
- **Event horizon**:  $\chi_{eh}(\tau) = \tau_f - \tau = \int_t^{\tau_f} \frac{dt}{a(t)}$  with  $t_f = \infty$

25

# SMC: Particle and Event horizons



**Figure 2.1:** Spacetime diagram illustrating the concept of horizons. Dotted lines show the worldlines of comoving objects. The event horizon is the maximal distance to which we can send signal. The particle horizon is the maximal distance from which we can receive signals.

# SMC: distances, angular sizes and volumes

- Comoving coordinate distance:

(also computed using photons that travel along null geodesics,  $ds^2 = 0$ , with  $d\theta = d\phi = 0$ )

$$ds^2 = c^2 dt^2 - a(t)^2 \frac{dr^2}{1 - kr^2} = 0 \quad \longrightarrow \quad \int_{r_0}^r \frac{dr}{\sqrt{1 - kr^2}} = c \int_{t_0}^t \frac{dt'}{a(t')}$$

- Proper (physical) distance:

$$d(t) = a(t) \int_{r_0}^r \frac{dr}{\sqrt{1 - kr^2}} \equiv \int_{r_0}^r \sqrt{|g_{rr}|} = a(t)c \int_{t_0}^t \frac{dt'}{a(t')}$$

From:

for a  $\Omega_\Lambda = 0$  universe this gives:

$$d(t) \simeq \frac{2}{3w + 1} \frac{c}{H_0} \Omega_{w0}^{1/2} \left( \frac{a}{a_0} \right)^{3(1+w)/2} = 3 \frac{1+w}{1+3w} ct$$

This equality holds only for  $\Omega_{w0} = 1$  (critical density universe,  $\Omega_k = \Omega_\Lambda = 0$ ). See Cosmology course notes)

27


# SMC: distances angular sizes and volumes

- Angular size of a region at a given time:

$$\theta = \frac{D}{d_A(t)}$$


where

$$d_A(t) = a(t) \int_{r_0}^r \frac{dr}{\sqrt{1 - kr^2}} \equiv \int_{r_0}^r \sqrt{|g_{rr}|} = a(t)c \int_{t_0}^t \frac{dt'}{a(t')}$$



**You can prove** that the angular size of the **particle horizon** at time/redshift  $z$  for a critical density with  $\Omega_\Lambda = 0$  is given by:

$$\theta_H \simeq 2 \tan \frac{\theta_H}{2} = \frac{\Omega_0^{3/2} \sqrt{1+z}}{\Omega_0 z + (\Omega_0 - 2) (\sqrt{1 + \Omega_0 z} - 1)}$$



# SMC: distances, angular sizes and volumes

- Hubble length:

Is defined as the length scale obtained when one sets  $v_H = c$  in the Hubble law  $v_H = c = H r$ .

$$R_H(t) = \frac{c}{H(t)} = \frac{3(w+1)}{2} ct$$

This equality holds only for  $\Omega_{w0} = 1$ , (critical density universe,  $\Omega_k = \Omega_\Lambda = 0$ ). See Cosmology course notes

- Physical volume element:

It is defined in the usual way “ $dV = dx dy dz$ ”. In spherical coordinates is:

$$dV = \sqrt{|g|} dr d\theta d\phi = (ar)^2 \frac{a dr}{\sqrt{1-kr^2}} d\Omega$$

You are also able to demonstrate that ( $d\Omega = d\theta d\phi$  is the solid angle)

$$\frac{dV}{d\Omega dz} = \frac{c}{H(z)} \frac{(a_0 r)^2}{(1+z)^3} = \frac{c}{H_0} \frac{d_A^2}{\mathcal{H}(z)(1+z)} \quad \text{where: } \mathcal{H}(z) = H(z)/H_0$$

29



Problems of the FLRW models  
as a sole ingredient of the SMC

# The Horizon Problem

At high redshift ( $z \gg 1$ ):

$$\theta_H \simeq \frac{180}{\pi} \sqrt{\frac{\Omega_0}{z}} \text{ deg}$$

At  $z_{cmb} \sim 1000$ ,  $\theta_H \simeq 1^\circ$  there are  $\sim 54000$  causal disconnected angular areas in the CMB sky. So, *why the CMB has a thermal spectrum with a so uniform temperature in all directions (2.725 °K)?*



# The Flatness Problem

From the Friedmann Equation, with  $\Lambda=0$ , one has

$$|\Omega(t) - 1| = \frac{|k|}{a^2(t)H^2(t)} = \frac{|k|}{\dot{a}^2(t)}$$

is a decreasing function of time:  
So as  $t \rightarrow 0$ ,  $\Omega \rightarrow 1$

**decreases tremendously** as time approaches the big bang instant.

This means that as we go back in time the **energy density of universe must be extremely close to the critical density** ( $t \rightarrow 0 \Rightarrow \Omega \rightarrow 1$ ). For  $t=1e-43$  s (Planck time)  $\Omega$  should deviate no more than  $1e-60$  from the unity!

*Why the universe has to “start” with  $\Omega(t)$  so close to 1?*<sup>32</sup>



# The Monopoles & other relics Problem

Particle physics predicts that a variety of **“exotic” stable particles**, such as the magnetic monopoles, should be produced in the early phase of the Universe and remain in measurable amounts until the present.

***No such particles have yet been observed. Why?***

This either implies that the predictions from particle physics are wrong, or their densities are very small and therefore there is something missing from this evolutionary picture of the Big Bang.




33

# The Origin of Perturbations Problem

Locally the universe is not homogeneous. It displays a complex hierarchical pattern of galaxies, clusters and super clusters.

*What's the origin of cosmological structure?  
Does it grow from gravitational instability?  
What is the origin of the initial perturbations?*


Without a mechanism to explain their existence one must assume that they “were born” with the universe already showing the correct amplitudes on all scales, so that gravity can accurately reproduce the present-day structures?




# The homogeneity and isotropy Problem

Why is the universe homogeneous on large scales? **At early times homogeneity had to be even more “perfect”.**

The **FLRW** universes form a **very special subset of solutions** of the GR equations. So, *why nature “prefers” homogeneity and isotropy from the beginning as opposed to having evolved into that stage?*



CMB T=2.725 K



The APM Galaxy Survey  
Maddox et al.



Distant Objects in the Hubble Ultra Deep Field  
NASA, ESA, R. Windhorst (Arizona State University) and H. Yan (Spitzer Science Center, Caltech)

## Theory of Inflation: solves the problems?

Inflation can be defined as

$$\text{Inflation} \Leftrightarrow \ddot{a} > 0 \Leftrightarrow \frac{d}{dt} (cH^{-1}/a) < 0.$$

This happens when

$$\left(\frac{\dot{a}}{a}\right)^2 = \frac{8\pi G}{3}\rho$$


$$\frac{\ddot{a}}{a} = -\frac{4\pi G}{3}\left(\rho + 3\frac{p}{c^2}\right) \quad \rightarrow \quad \ddot{a} > 0 \Leftrightarrow \rho + \frac{3p}{c^2} < 0 \Leftrightarrow$$

$$p < -\rho c^2/3$$


**Riddle:** no known matter / energy component has an equation of state parameter  $w = \rho c^2/p < -1/3...$   
(continues in Chapter 9)

Appendix 1  
CMB angular power spectrum:  
a primer

Why an “elliptical” map?



# Earth's "elliptical" map (mollweide projection)



## CMB: temperature fluctuations on the sphere

- Can be expanded as a sum of functions, the spherical harmonics  $Y_{lm}$ , that are a basis on the surface of a sphere:

$$\Theta(\hat{n}) = \Delta T/T(\theta, \phi) = \sum_{\ell=0}^{\infty} \sum_{m=-\ell}^{\ell} a_{\ell m} Y_{\ell m}(\theta, \phi)$$

- The coefficients  $a_{\ell m}$  are the projection of the temperature fluctuation function onto the basis function  $Y_{\ell m}$  (it measures the contribution of a given  $Y_{\ell m}$  function to the temperature fluctuation):

$$a_{\ell m} = \int Y_{\ell m}^*(\theta', \phi') \frac{\Delta T}{T}(\theta', \phi') d\Omega'$$


- The angular power spectrum is defined as an angular correlation function in the celestial sphere:

$$C(\hat{n}, \hat{n}') \equiv \left\langle \frac{\Delta T}{T}(\hat{n}) \frac{\Delta T}{T}(\hat{n}') \right\rangle = \sum_{\ell \ell'} \sum_{m m'} (a_{\ell m}^* a_{\ell' m'}) Y_{\ell m}^*(\hat{n}) Y_{\ell' m'}(\hat{n}')$$


C<sub>l</sub>

# CMB angular power spectrum

Planck




# CMB angular power spectrum




# CMB angular power spectrum

Figure credits: Wayne Hu




# Spherical harmonics



# Spherical harmonic components of a well know map...

$\ell=1$


$$\sum_m a_{1m} Y_{1m}$$



Made by Matthias Bartelmann

$\ell=2$


$$\sum_m a_{2m} Y_{2m}$$



Made by Matthias Bartelmann

$\ell=3$


$$\sum_m a_{3m} Y_{3m}$$



Made by Matthias Bartelmann

$\ell=4$

$$\sum_m a_{4m} Y_{4m}$$




Made by Matthias Bartelmann



$\ell=5$


$$\sum_m a_{5m} Y_{5m}$$



Made by Matthias Bartelmann

$\ell=6$


$$\sum_m a_{6m} Y_{6m}$$



Made by Matthias Bartelmann

$\ell=7$

$$\sum_m a_{7m} Y_{7m}$$



Made by Matthias Bartelmann


$\ell=8$

i.e. larger  $\ell$  means shorter wavelengths

i.e.  $\ell$  is spherical equivalent of wavenumber

$\ell \sim \pi / \theta$


$$\sum_m a_{8m} Y_{8m}$$



Made by Matthias Bartelmann

$\ell=1$


$$\sum_{l=1}^1 \sum_m a_{lm} Y_{lm}$$



Made by Matthias Bartelmann

$\ell=1$  plus  $\ell=2$


$$\sum_{l=1}^2 \sum_m a_{lm} Y_{lm}$$



Made by Matthias Bartelmann

$\ell=1$  plus  $\ell=2$  plus  $\ell=3$


$$\sum_{l=1}^3 \sum_m a_{lm} Y_{lm}$$



Made by Matthias Bartelmann

Sum  $\ell=1$  to 4


$$\sum_{l=1}^4 \sum_m a_{lm} Y_{lm}$$



Made by Matthias Bartelmann

Sum  $\ell=1$  to 5


$$\sum_{l=1}^5 \sum_m a_{lm} Y_{lm}$$



Made by Matthias Bartelmann

Sum  $\ell=1$  to 6


$$\sum_{l=1}^6 \sum_m a_{lm} Y_{lm}$$



Made by Matthias Bartelmann

Sum  $\ell=1$  to 7


$$\sum_{l=1}^7 \sum_m a_{lm} Y_{lm}$$



Made by Matthias Bartelmann

Sum  $\ell=1$  to 8


$$\sum_{l=1}^8 \sum_m a_{lm} Y_{lm}$$



Made by Matthias Bartelmann

Sum up to some high  $\ell$


$$\sum_{l=1}^{100} \sum_m a_{lm} Y_{lm}$$




Made by Matthias Bartelmann

Earth's map with all contributions up to Planck's CMB map resolution

$$\sum_{l=1}^{1000} \sum_m a_{lm} Y_{lm}$$



# CMB temperature fluctuations map



Goal:

Use CMB map information to constrain theoretical Cosmological models

## Online $C_l$ calculators

NASA National Aeronautics and Space Administration

RSS LAMBDA News

Search / Site Map

+ HOME + PRODUCTS - TOOLBOX + LINKS + NEWS + SITE INFO

LEGACY ARCHIVE FOR MICROWAVE BACKGROUND DATA ANALYSIS

"One Stop Shopping for CMB Researchers"

**CMB Toolbox**

- + Tools
- + Contributed S/W
- + CAMB
  - Online Tool
    - + Overview
    - + CMBFAST
      - + Online Tool
      - + Overview
    - + WMAPViewer
      - + Online Tool
      - + Overview
    - + Conversion Utilities

**CAMB Web Interface**

Supports the September 2008 Release

Most of the configuration documentation is provided in the sample parameter file provided with the application.

This form uses JavaScript to enable certain layout features, and it uses Cascading Style Sheets to control the layout of all the form components. If either of these features are not supported or enabled by your browser, this form will NOT display correctly.

Actions to Perform

Scalar  $C_l$ 's  Do Lensing  Linear

Vector  $C_l$ 's  Transfer Functions  Non-linear Matter Power (HALOFIT)

Tensor  $C_l$ 's  Non-linear CMB Lensing (HALOFIT)

Sky Map Output:

Vector  $C_l$ 's are incompatible with Scalar and Tensor  $C_l$ 's. The Transfer functions require Scalar and/or Tensor  $C_l$ 's.

The HEALpix synfast program is used to generate maps from the resultant spectra. The random number seed governs the phase of the  $a_m$ 's generated by synfast. The default of zero causes synfast to generate a new seed from the system time with each run. Specifying a fixed nonzero value will return fixed phases with.

CMB Toolbox: <http://lambda.gsfc.nasa.gov/toolbox/>


CAMB website: <http://camb.info/>

CMBFast website: <http://www.cmbfast.org/>



# CMB analyzer

[http://lambda.gsfc.nasa.gov/education/cmb\\_plotter/](http://lambda.gsfc.nasa.gov/education/cmb_plotter/)



This tool requires an Adobe Flash player to be installed for your browser.  
Make full screen in new window

35

## CMB parameter cheat sheet

