

# Universo Primitivo

## 2023-2024 (1º Semestre)

Mestrado em Física - Astronomia


### Chapter 1

#### 1. The observed Universe

- Foreword: The Olbers' paradox;
- The Universe at different scales;
- Observational Cosmology: empirical facts and the hot Big-Bang theory
  - Cosmic Expansion: The Hubble law;
  - The abundancies of the light elements;
  - The existence of a Cosmic Background Radiation;
  - The isotropy of distant objects;
  - The existent of dark matter;
  - The accelerated expansion of the Universe
- Formation and evolution of cosmic structure

# Foreword: The Olbers' paradox and the present view of the Universe


## Foreword: Why is the sky dark at night?



Heinrich Olbers  
(1758–1840)



# Foreword: Why is the sky dark at night?




Heinrich Olbers  
(1758–1840)

**Olbers' paradox** (1826) : argues that “the **darkness of the sky** at night **conflicts with the concept of an infinite and eternal static universe**”, with stars distributed uniformly.

$$\text{light} = \int_{r_0}^{\infty} L(r)N(r) dr$$



# Foreword: Why is the sky dark at night?




Heinrich Olbers  
(1758–1840)


**Olbers' paradox** (1826) : argues that “the **darkness of the sky** at night **conflicts with the concept of an infinite and eternal static universe**”, with stars distributed uniformly.

Olbers paradox in action.

Exercise: prove why this happens




# Foreword: Why is the sky dark at night?



Heinrich Olber's  
(1758–1840)


Some possible explanations:

1. Too much dust absorbs light from distant stars.
2. The number of stars in the Universe is finite.
3. The distribution of stars is not uniform.
4. The Universe is expanding. Light from distant stars are dimmed (redshifted) into obscurity.
5. The observed Universe has a finite age. Distant light hasn't even reached us yet.



The Universe at different  
scales

## Stars...



Sun: 8 light minutes away  
 $\alpha$ -Centauri 4,25 light years

## Gaseous Nebulae...




Distance to the Eagle Nebulae 7000 light years


The galaxy...



A home for billions of stars...





# Milky Way



16 000 light years


120 000 light years

# Groups and Clusters of Galaxies...




Distance to Coma cluster 321 million ly – 99 Mpc

# Our place in the Universe...



Credits: American Natural History Museum; gently provided by Miguel de Avillez U. Évora

# LSS: “The fingers of God”...



*Credit: M. Blanton and the Sloan Digital Sky Survey*

Further away and back in time...  
First stars forming period

Distance: 12.8 – 13.4 billion years back in time



CMB...

The edge of the visible Universe


Distance: ~13.8 billion years back in time

Fig. credits: NASA / WMAP Science Team

**Planck Surveyor:** looking back to the dawn of time



BR-275



**Project:** ESA lead mission to observe the temperature and polarization anisotropies of the Cosmic Microwave Background (CMB) radiation with unprecedented precision.

**Total Cost:** about €700 million (€1 / person in EU)

**Mission timeline:**

**Launch:** 14 May 2009

**Operational orbit at L2:** July 2009

**Nominal science phase:** end of January 2011

**Extended mission:** Shut down date: 19 Oct. 2013

**Payload:**

**Telescope:** 1.5 m projected apertures


**Low Frequency Instrument (LFI):** array of 22 tuned radio receivers operating at 30, 44 and 70 GHz.

**High Frequency Instrument (HFI):** array of 52 bolometers operating at 100, 143, 217, 353, 545, and 857 GHz.

Fig. credits: ESA

# Planck CMB observations


2009-2013: Planck satellite observes the CMB sky with unprecedented angular resolution and sensitivity.



Animation credits: ESA and the Planck collaboration; Cluster map by Douspis, Hurier, Aghanim 2013

# Planck CMB observations

2009-2013: Planck satellite observes the CMB sky with unprecedented angular resolution and sensitivity.




Animation & Fig. credits: ESA and the Planck collaboration

# Galaxy surveys: 3D mapping of the Universe...


SDSS: aims at ~25% of the sky; ~100 million objects

## ESA Cosmology missions




ESA/NASA Planck Surveyor:  
2009 - 2013

ESA Euclid:  
2023 - 2029



# Euclid mission (ESA): Galaxy Surveys from space (launch 2023)

Portuguese official participation lead by the cosmology group @ IA – CAUP/FCUL



Euclid: ~ 2000 million galaxies

Fig. credits: ESA - C. Carreau.

# Euclid mission (ESA): Galaxy Surveys from space (launch 2023)

**Primary Objective:** To map the distribution and evolution of (dark) matter in the Universe to:

- Measure the accelerated expansion
- Unveil the nature of dark energy
- Test models of gravity, dark matter, and of the primordial universe

Present observations suggest that dark energy (DE) and dark matter (DM) are 96% of the total energy density budget of the universe.






Fig. credits: ESA - C. Carreau.

## 3D galaxy surveys



SDSS DR3




SDSS DR4

Subbarao, Surendran, e Landsberg. (SDSS)

## Cartography of galaxies in 3D with Euclid




O Euclid fará a maior e mais precisa cartografia, de galáxias no Universo, observando imagens e obtendo espectros a partir do espaço.




# Cartography of galaxies in 3D

SDSS DR3



cosmicweb.barabasilab.com




Subbarao, Surendran, e Landsberg. (SDSS)

Teia cósmica


# Analogy with the study of living organisms

Cosmic web



cosmicweb.barabasilab.com



Spider web



# Spider webs (effect of caffeine concentration)





# Cosmic webs (with different amounts of DM & DE)




# Cosmic web probes cosmology & particle physics parameters

Model	$\Omega_{m0}$	$\lambda$	$h$	$\Gamma$	$\Sigma_8$	$N_{par}$	$L$ (Mpc/h)	$m_p$ ( $M_{sun}/h$ )	$L_{soft}$ (Kpc/h)
SCDM	1.0	0.0	0.5	0.50	0.6	$256^3$	84.5	$1.0 \times 10^{10}$	36
$t$ CDM	1.0	0.0	0.5	0.21	0.6	$256^3$	84.5	$1.0 \times 10^{10}$	36
$\Lambda$ CDM	0.3	0.7	0.7	0.21	0.90	$256^3$	141.3	$1.4 \times 10^{10}$	20
OCDM	0.3	0.0	0.7	0.21	0.85	$256^3$	141.3	$1.4 \times 10^{10}$	30




← Impact of Dark Matter & Dark Energy densities; Hubble constant; Primordial power spectrum

Impact of the nature of the Dark Matter (Part. Physics model) →



## Euclid: an M-class mission of ESA's scientific program

Euclid will observe more than a billion of galaxies during 6 years of operations. The light emitted by galaxies is lensed (deflected) by invisible structures of Dark Matter and are observed with distortion.



## Euclid: an M-class mission of ESA's scientific program

The distortions are very small. But with the observation of a billion galaxies, there is enough statistics to map the dark matter and study the nature of dark energy.




Imagem do Hubble do enxame de galáxias Abell 370  
(a 4 bilhões de anos luz da terra)




# Euclid: an M-class mission of ESA's scientific program

To obtain good images, the satellite will be placed in orbit at the Lagrange point L2, with its "back turned" to the sun, pointing successively to different directions of the darkest regions of the sky. Each field will have to be observed for 1h15. The total area of observation will be 15000 square degrees.



# Euclid Sky Survey implementation and optimization is a PT, national, responsibility

<https://www.youtube.com/watch?v=y89SXN-Xd9A>




Animação criada por J. Dinis (FCUL / IA)


# Euclid Survey planning

Portuguese official contribution is carried out at IA/FCUL:

- I. Tereno
- J. Dinis
- C. S. Cavalho
- A. da Silva



12 de outubro de 2023



From: Euclid Preparation I. The Euclid Reference Survey: status at the Preliminary Design Review, submitted to the ECEB, 2019

DF-FCUL, Lisboa

35

Observational cosmology:  
empirical facts about the  
Universe

# 1. The Universe is expanding






Edwin Hubble

1912: Vesto Slipher is the first to observe spectral line (red)shifts towards nebulae and to relate these redshifts to their recessional velocities.

1924: Edwin Hubble ends debate on the nature of nebulae being galactic objects

1929: reports a linear relation between relative radial velocity and distance:  $v = Hd$



From:

# 1. The Universe is expanding




Edwin Hubble


1924: Edwin Hubble ends debate on the nature of nebulae being galactic objects

1929: reports a linear relation between relative radial velocity and distance:  $v = Hd$

At a given time



At a latter time, t



From:

# 1. The Universe is expanding




Edwin Hubble


1924: Edwin Hubble ends debate on the nature of nebulae being galactic objects

1929: reports a linear relation between relative radial velocity and distance:  $v = Hd$

Time evolution of an expanding spherical surface



From:



## The basic idea behind the Big-Bang theory

- If the universe is expanding and matter-energy is conserved during the expansion then the universe had to be smaller, denser and hotter in the past!
- If so, the Universe must have evolved from a state where matter and radiation form a **ultra dense and hot ionized plasma** of fundamental particles
- As the universe expands and cools down:
  - interactions between the plasma components become less frequent;
  - different particle species should decouple from the plasma;
  - eventually the universe becomes neutral and transparent to radiation



# According to the Big-Bang theory, in the early instants...

*“ the Universe was a extremely hot and dense plasma, like a ‘torrid bright fog’ ...*

*... radiation was trapped in this plasma through collisions with other plasma particles*

*... as the universe expands, the plasma temperature drops, atomic nuclei form, and capture the free electrons in the plasma. When the number of free electrons is too small, radiation no longer interacts with the plasma and propagates freely, giving rise to the Cosmic Microwave Background and neutral matter”*

41

## 2. The abundance of light nuclei




Herman, Gamow, Alpher

The relative abundance of light elements can not be explained by stellar nucleosynthesis

1948: Alpher & Gamow computed the abundance of light elements in the context of the **Big Bang** theory

Light elements were produced at low temperatures ( $< 10^9$ K and high densities) during several tens of minutes



From:



# 3. Cosmic Microwave Background




Penzias & Wilson

1965: Penzias & Wilson serendipitously discovered a uniform radiation (“excess”) across the sky.


This was the cosmic microwave background radiation predicted by Gamow and Alpher in 1948



## CMB: the last scattering surface





# 3. Cosmic Microwave Background



John Mather & George Smoot

1991: High precision measurement of CMB temperature by COBE and 1st detection of temperature fluctuations (Mather & Smoot)

2001: State of the art measurements of  $dT/T \sim 1e-5$  temperature fluctuations by WMAP




WMAP

Fig. credits: NASA / WMAP Science Team

# 4. Isotropy of distant objects


On Large Scales the Universe...  
... appears to be ISOTROPIC

+




The Cosmological Principle (Milne, Einstein):  
"all places in the Universe are alike" — Einstein

The Universe Homogeneous and Isotropic on large scales



The APM Galaxy Survey  
Maddox et al



Distant Objects in the Hubble Ultra Deep Field  
NASA, ESA, R. Windhorst (Arizona State University) and H. Yan (Spitzer Science Center, Caltech)

# 4. Isotropy of distant objects






Fig. 3.1 A point distribution, statistically isotropic around every point (left) and around a unique point ( $P$ ) (right). In the second version,  $P$  and  $Q$  are not equivalent. The cosmological principle excludes such kinds of solutions, which would assume that we lie in a special place in the Universe. From Ref. [1] of the introduction.




The APM Galaxy Survey  
Maddox et al

NASA, ESA, R. Windhorst (Arizona State University) and H. Yan (Spitzer Science Center, Caltech)

# 5. The existence of Dark Matter



From:




Jan Oort

1927: Jan Oort studies the **rotation of stars in our galaxy** and infers that their rotation is not consistent with Keplerian motion.

$$v_{circ} = \sqrt{\frac{GM(R)}{r}}$$




# 5. The existence of Dark Matter



Oorts constants:

$$A \equiv -\frac{1}{2} \left[ \frac{dV_c}{dR} \Big|_{R_0} - \frac{V_{c,0}}{R_0} \right]$$

$$B \equiv -\frac{1}{2} \left[ \frac{dV_c}{dR} \Big|_{R_0} + \frac{V_{c,0}}{R_0} \right]$$



Circular motion:

$$v_{circ} = \sqrt{\frac{GM(R)}{r}}$$

If the whole mass is mostly at the centre:  $v_{cir}^2 \sim 1/r$


Observations vs Keplerian motion:

- Kepler. motion:  $(A-B)/(A+B) = 2$
- Observations :  $(A-B)/(A+B) = 5$

-Mass is not concentrated at the centre  
-Non-luminous mass is required

<http://icc.dur.ac.uk/~tt/Lectures/Galaxies/TeX/lec/node42.html>

# 5. The existence of Dark Matter



B: Observations

A: theoretical expectations

1980: Vera Rubin and others also find that **stars rotate too fast in the outskirts of spiral galaxies** to remain bound assuming that gravity is produced only by visible matter.

# 5. The existence of Dark Matter




Fritz Zwicky

1936: Fritz Zwicky applied the **Virial theorem to the velocities of galaxies in the Coma cluster** and finds very high mass-to-light ratios,  $\Upsilon = M/L$ , for them to remain bound:  $\Upsilon_{coma}/\Upsilon_{sun} = 500 \gg 2-10$  for galaxies.


- **Virial theorem (for gravitationally relaxed systems):**  $2\bar{E}_k + \bar{E}_p = 0$
- **Mass from the virial theorem:**  $M_V = \langle v^2 \rangle R / G$
- **Visible luminous Mass:**  $M_L = N_g \Upsilon_g L_g$   
( $N_g$  - number of galaxies;  $\Upsilon_g$  - galaxy mass-to-light ratio;  $L_g$  galaxy luminosity)

# 5. The existence of Dark Matter


lensing effects:



Albert Einstein




$$G_{\mu\nu} = 8\pi G T_{\mu\nu}$$



From:


# 5. The existence of Dark Matter

lensing effects:



From:


**Strong lensing**



# 5. The existence of Dark Matter

lensing effects: **strong lensing**

**Einstein Rings**




Galaxy-galaxy lensing





Double Einstein Ring SDSSJ0946+1006

Hubble Space Telescope • ACS/WFC



# 5. The existence of Dark Mater

2003: X-ray (produced by extremely hot gas – in red) vs weak lensing observations (probing the total mass distribution in blue) of the **Bullet Cluster** put in evidence that galaxy clusters must contain “dark matter”



Refs:  
[astro-ph/0309303](https://arxiv.org/abs/astro-ph/0309303)  
[astro-ph/0312273](https://arxiv.org/abs/astro-ph/0312273)

# 6. Cosmic expansion is accelerating




1998: S. Perlmutter and the supernova Cosmology project found first evidence for the accelerated expansion of the Universe.

assuming supernovae are standard candles, they appear further away (green arrow) than predicted by non-accelerating expansion models (yellow arrow).


$$d(t) = a(t) d_0$$

with  $\ddot{a}(t) > 0$

No Accelerated expansion






Accelerated expansion




From:

# 6. Cosmic expansion is accelerating




Cosmological redshift:

$$\begin{aligned}
 z &= \frac{E - E_0}{E_0} = \\
 &= \frac{\nu}{\nu_0} - 1 = \\
 &= \frac{\lambda_0}{\lambda} - 1 = \\
 &= \frac{a}{a_0} - 1
 \end{aligned}$$



# How Cosmic structure forms and evolves?





## Observations indicate that

- ❑ on small scales the universe is NOT homogeneous and isotropic
- ❑ On large cosmological scales the Universe does not show indications of strong anisotropies. Together with the cosmological principle this implies the universe is highly homogeneous and isotropic
- ❑ However it shows small anisotropies in the CMB.

# How structure forms and evolves?

Structures should grow due to  
Gravitational Instability (Jeans, Lifshitz,...)



$$\mathbf{g}(\mathbf{r}, t) = -\frac{1}{a} \nabla \phi = \frac{3\Omega H^2}{8\pi} \int d\mathbf{x}' \delta(\mathbf{x}', t) \frac{(\mathbf{x}' - \mathbf{x})}{|\mathbf{x}' - \mathbf{x}|^3}$$

Density fluctuations:  $t=13.7$  billion years

$$\mathbf{g}(\mathbf{r}, t) = -\frac{1}{a} \nabla \phi = \frac{3\Omega H^2}{8\pi} \int d\mathbf{x}' \delta(\mathbf{x}', t) \frac{(\mathbf{x}' - \mathbf{x})}{|\mathbf{x}' - \mathbf{x}|^3}$$

## Density fluctuations: t=13.5 billion years

500 Mpc/h

$$\mathbf{g}(\mathbf{r}, t) = -\frac{1}{a} \nabla \phi = \frac{3\Omega H^2}{8\pi} \int d\mathbf{x}' \delta(\mathbf{x}', t) \frac{(\mathbf{x}' - \mathbf{x})}{|\mathbf{x}' - \mathbf{x}|^3}$$

Millennium simulation, Springel et al.

## Density fluctuations: t=12.7 billion years

500 Mpc/h

$$\mathbf{g}(\mathbf{r}, t) = -\frac{1}{a} \nabla \phi = \frac{3\Omega H^2}{8\pi} \int d\mathbf{x}' \delta(\mathbf{x}', t) \frac{(\mathbf{x}' - \mathbf{x})}{|\mathbf{x}' - \mathbf{x}|^3}$$

Millennium simulation, Springel et al.

## Density fluctuations: t=9 billion years

500 Mpc/h

$$\mathbf{g}(\mathbf{r}, t) = -\frac{1}{a} \nabla \phi = \frac{3\Omega H^2}{8\pi} \int d\mathbf{x}' \delta(\mathbf{x}', t) \frac{(\mathbf{x}' - \mathbf{x})}{|\mathbf{x}' - \mathbf{x}|^3}$$

Millennium simulation, Springel et al.

## Density fluctuations: t=1 billion years

500 Mpc/h

$$\mathbf{g}(\mathbf{r}, t) = -\frac{1}{a} \nabla \phi = \frac{3\Omega H^2}{8\pi} \int d\mathbf{x}' \delta(\mathbf{x}', t) \frac{(\mathbf{x}' - \mathbf{x})}{|\mathbf{x}' - \mathbf{x}|^3}$$

Millennium simulation, Springel et al.



# Cosmological structure formation

$z = 20.0$

50 Mpc/h



# Large Scale Structure (LSS)

1 Gpc/h




Millennium Simulation

10,077,696,000 particles

( $z = 0$ )

# Large Scale Structure (LSS)



# The history of the Universe:

