

Universo Primitivo

2018-2019 (1º Semestre)


Mestrado em Física - Astronomia

Chapter 10

10 Inflation: the origin of perturbations

- The Basic Picture;
- Cosmological perturbation theory
- Quantum fluctuations in the de Sitter space;
- Primordial power spectra from inflation;
- CMB power spectrum

References



Inflation: the basic picture

The Inflationary phase of the Universe would need to happen at very early times. Present data is consistent with an inflationary period that lasted for about around $\Delta t \sim 10^{-36}$ at cosmic time of about $t \sim 10^{-32} - 10^{-33}$ seconds

In these conditions the **inflaton field has a quantum nature** and its energy density is quantified. The **Heisenberg uncertainty principle** allows the origin of energy density fluctuations given the short timescales involved.

$$\Delta E_\phi > h/(4\pi\Delta t)$$

The **inflation field**, $\phi(x, t)$, therefore **acquires a spatial dependence due to quantum fluctuations**, $\delta\phi(x, t)$, about its “background” value, $\phi(t)$:

$$\phi(x, t) = \phi(t) + \delta\phi(x, t)$$




Figure 6.1: Quantum fluctuations $\delta\phi(t, x)$ around the classical background evolution $\bar{\phi}(t)$. Regions acquiring a negative fluctuations $\delta\phi$ remain potential-dominated longer than regions with positive $\delta\phi$. Different parts of the universe therefore undergo slightly different evolutions. After inflation, this induces density fluctuations $\delta\rho(t, x)$.

Relativistic (GR) perturbation theory

Scalar, vector Tensor (SVT) decomposition

The perturbation variables can be decomposed into their scalar, vector and tensor dependences. This is useful because these dependences do not mix at linear order:

$$B_i = \underbrace{\partial_i B}_{\text{scalar}} + \underbrace{\hat{B}_i}_{\text{vector}}$$

$$h_{ij} = \underbrace{2C\delta_{ij} + 2\partial_{(i}\partial_{j)}E}_{\text{scalar}} + \underbrace{2\partial_{(i}\hat{E}_{j)}}_{\text{vector}} + \underbrace{2\hat{E}_{ij}}_{\text{tensor}}$$

with,

$$\partial_{(i}\partial_{j)}E \equiv \left(\partial_i\partial_j - \frac{1}{3}\delta_{ij}\nabla^2\right)E,$$

$$\partial_{(i}\hat{E}_{j)} \equiv \frac{1}{2}\left(\partial_i\hat{E}_j + \partial_j\hat{E}_i\right).$$

where:

SVT d.o.f.	{	4	• scalars: A, B, C, E		
		4	• vectors: \hat{B}_i, \hat{E}_i		$\partial^i \hat{B}_i = 0$
		2	• tensors: \hat{E}_{ij}		$\partial^i \hat{E}_i = 0$ and $\partial^i \hat{E}_{ij} = 0$

Relativistic (GR) perturbation theory

Gauge freedom

GR is a gauge theory where the gauge transformations are generic coordinate transformations.

$$ds^2 = g_{\mu\nu}(X)dX^\mu dX^\nu = \tilde{g}_{\alpha\beta}(\tilde{X})d\tilde{X}^\alpha d\tilde{X}^\beta$$



$$g_{\mu\nu}(X) = \frac{\partial \tilde{X}^\alpha}{\partial X^\mu} \frac{\partial \tilde{X}^\beta}{\partial X^\nu} \tilde{g}_{\alpha\beta}(\tilde{X})$$

A gauge choice is a way of choosing the (time) slicing and (spatial) threading of spacetime.

GAUGE CHOICE \iff SLICING AND THREADING

How to treat Perturbations?

- Either find gauge invariant variables to describe perturbations. These variables are called real spacetime perturbations.
- Or fix a gauge choice and keep track of all perturbations and check how quantities transform.



Relativistic (GR) perturbation theory

Gauge-invariant perturbation variables

One avoids gauge problems by defining special combinations of the SVT perturbations that do not change under coordinate transformations. These are known as the **Bardeen potentials** (or Bardeen Variables)

$$\begin{aligned}\Psi &\equiv A + \mathcal{H}(B - E') + (B - E')', & \hat{\Phi}_i &\equiv \hat{E}'_i - \hat{B}_i, & \hat{E}_{ij} \\ \Phi &\equiv -C - \mathcal{H}(B - E') + \frac{1}{3}\nabla^2 E.\end{aligned}$$

where $'$ is derivative with respect to conformal time, τ , and $\mathcal{H} \equiv a'/a$ is the Hubble parameter in conformal time.

Useful Gauge fixing choices

The gauge freedom can be used to conveniently set some of the above variables to zero:

- **Newtonian Gauge:** $E = B = 0$

The metric simply becomes:

$$ds^2 = a^2(\tau) [(1 + 2\Psi)d\tau^2 - (1 - 2\Phi)\delta_{ij}dx^i dx^j]$$

where the remaining non-zero variables were renamed to $A \equiv \Psi$, $C \equiv -\Phi$

Relativistic (GR) perturbation theory

Useful Gauge fixing choices

(continuation)

- **Spatially flat gauge** : $C = E = 0$

This is a convenient gauge choice for the calculation of the inflationary perturbations.

- **Uniform density gauge:** consists in choosing the time-slicing in a way that the total density perturbation (see perturbed stress-energy tensor subsection) is set to zero: $\delta\rho = 0$

- **Comoving gauge:** consists in choosing coordinates in a way that the total momentum density vanishes (see perturbed stress-energy tensor subsection): $q_i = (\bar{\rho} + \bar{P})v_i = 0$. One has that $q_i = B_i = 0$.

This choice is naturally connected to the inflationary initial conditions

Relativistic (GR) perturbation theory

Perturbed Stress-Energy Tensor

For small perturbations the perturbed stress-energy tensor can be written as:

$$T^\mu{}_\nu = \bar{T}^\mu{}_\nu + \delta T^\mu{}_\nu$$

where the unperturbed stress-energy tensor is

$$\bar{T}^\mu{}_\nu = (\bar{\rho} + \bar{P})\bar{U}^\mu\bar{U}_\nu - \bar{P}\delta^\mu{}_\nu$$

and one has that, $\bar{U}_\mu = a\delta_\mu^0$, $\bar{U}^\mu = a^{-1}\delta_0^\mu$, for a comoving observer.

The perturbation to the stress-energy tensor can be written as:

$$\delta T^\mu{}_\nu = (\delta\rho + \delta P)\bar{U}^\mu\bar{U}_\nu + (\bar{\rho} + \bar{P})(\delta U^\mu\bar{U}_\nu + \bar{U}^\mu\delta U_\nu) - \delta P\delta^\mu{}_\nu - \Pi^\mu{}_\nu$$

where $\Pi^\mu{}_\nu$ is the anisotropic stress tensor and the perturbed density, pressure and four-velocity vectors generally depend on space and time.

To 1st order one has (see eg Baumann):

$$\delta U^\mu = a^{-1}[-A, v^i]; \quad \delta U_\nu = a[A, -(v^i + B_i)]$$

and

$$U^\mu = a^{-1}[1 - A, v^i]; \quad U_\nu = a[1 + A, -(v^i + B_i)]$$

11

Relativistic (GR) perturbation theory

Perturbed Stress-Energy Tensor

(continuation)

Using these expressions of U^μ and U_ν in δU^μ one gets

$$\begin{aligned} \delta T^0{}_0 &= \delta\rho, \\ \delta T^i{}_0 &= (\bar{\rho} + \bar{P})v^i, \\ \delta T^0{}_j &= -(\bar{\rho} + \bar{P})(v_j + B_j), \\ \delta T^i{}_j &= -\delta P\delta_j^i - \Pi^i{}_j. \end{aligned}$$

The quantity $q_i = (\bar{\rho} + \bar{P})v_i$ is called the **momentum density three-vector**. Note that the perturbed (peculiar) velocity $\delta U^i \equiv v^i/a$ is not additive quantity, but q_i is additive. If there are several fluid components all the quantities bellow are additive:

$$\delta\rho = \sum_I \delta\rho_I, \quad \delta P = \sum_I \delta P_I, \quad q^i = \sum_I q^i_I, \quad \Pi^{ij} = \sum_I \Pi^{ij}_I$$

And the stress-energy tensor is also additive: $T_{\mu\nu} = \sum_I T_{\mu\nu}^I$

The **SVT decomposition** can also be applied to the perturbed stress-energy tensor: $\delta\rho$ and δP only have scalar parts; $q_i = \partial_i q + \hat{q}_i$ has a scalar and a vector part; Π_{ij} has scalar, vector and tensor parts: $\Pi_{ij} = \partial_{\langle i}\partial_{j\rangle}\Pi + \partial_{(i}\hat{\Pi}_{j)} + \hat{\Pi}_{ij}$

12

Relativistic (GR) perturbation theory

Gauge-invariant perturbation quantities

Comoving-gauge density perturbation: The quantity :

$$\bar{\rho}\Delta \equiv \delta\rho + \bar{\rho}'(v + B)$$

Where v is a scalar velocity function such that $v_i = \partial_i v$, is gauge-invariant. It is very useful to study density perturbations .

Comoving Curvature perturbation: In a arbitrary gauge, the intrinsic curvature of hypersurfaces of constant time can be computed using the special part of the perturbed metric. Since this is a scalar it only receives contributions from the scalar variables of the spatial part of metric ($E_{ij} \equiv \partial_{(i}\partial_{j)}E$) :

$$\gamma_{ij} \equiv a^2 [(1 + 2C)\delta_{ij} + 2E_{ij}]$$

After some long calculations (see Baumann) the intrinsic curvature is given by:

$$a^2 R_{(3)} = -4\nabla^2 \left(C - \frac{1}{3}\nabla^2 E \right)$$

The comoving curvature perturbation

$$\mathcal{R} = C - \frac{1}{3}\nabla^2 E + \mathcal{H}(B + v)$$

Is gauge-invariant and it is defined as the comoving curvature computed in the comoving gauge ($q_i = B_i = 0$). In the Newtonian gauge this is $\mathcal{R} = -\Phi + \mathcal{H}v$.

13

Relativistic (GR) perturbation theory

Adiabatic versus Isocurvature perturbations

Density perturbations are said to be **adiabatic** if

$$\delta\rho_I(\tau, \mathbf{x}) \equiv \bar{\rho}_I(\tau + \delta\tau(\mathbf{x})) - \bar{\rho}_I(\tau) = \bar{\rho}'_I \delta\tau(\mathbf{x})$$

for all fluid components, I . This implies:

$$\delta\tau = \frac{\delta\rho_I}{\bar{\rho}'_I} = \frac{\delta\rho_J}{\bar{\rho}'_J} \quad \text{for all species } I \text{ and } J$$

If fluid components obey to independent continuity equations, $\bar{\rho}'_I = -3\mathcal{H}(1 + w_I)\bar{\rho}_I$ one gets:

$$\frac{\delta_I}{1 + w_I} = \frac{\delta_J}{1 + w_J} \quad \text{for all species } I \text{ and } J$$

This also implies that the total density density of the fluid is perturbed and is given simply by

$$\delta\rho_{\text{tot}} = \bar{\rho}_{\text{tot}}\delta_{\text{tot}} = \sum_I \bar{\rho}_I \delta_I$$

14

Relativistic (GR) perturbation theory

Adiabatic versus Isocurvature perturbations

(continuation)

Isocurvature perturbations are perturbation in the different fluid components in a way that conserves the total energy density. This implies that different fluid components have fluctuations such as the quantity:

$$S_{IJ} \equiv \frac{\delta_I}{1 + w_I} - \frac{\delta_J}{1 + w_J}$$

is different from zero.

Linear perturbation GR equations & conservation laws

Once the perturbed stress-energy tensor and perturbed metric are defined one proceeds with the calculation of the:

- Perturbed metric connections;
- The conservation laws of the perturbed stress-energy tensor;
- The Einstein equations involving the perturbed quantities up to linear order of the perturbed quantities (higher order calculations are more complex or impossible to do). (e.g. Ch.4 Baumann)
- Solve the resulting equations to derive the evolution of perturbations (e.g. Ch.5 Baumann)

15

Relativistic (GR) perturbation theory

Linear perturbation GR equations & conservation laws (Newton. gauge)

$$ds^2 = a^2(\tau) [(1 + 2\Psi)d\tau^2 - (1 - 2\Phi)\delta_{ij}dx^i dx^j] . \quad (4.4.168)$$

In these lectures, we won't encounter situations where anisotropic stress plays a significant role, so we will always be able to set $\Psi = \Phi$.

- The Einstein equations then are

$$\nabla^2\Phi - 3\mathcal{H}(\Phi' + \mathcal{H}\Phi) = 4\pi G a^2 \delta\rho , \quad (4.4.169)$$

$$\Phi' + \mathcal{H}\Phi = -4\pi G a^2 (\bar{\rho} + \bar{P})v , \quad (4.4.170)$$

$$\Phi'' + 3\mathcal{H}\Phi' + (2\mathcal{H}' + \mathcal{H}^2)\Phi = 4\pi G a^2 \delta P . \quad (4.4.171)$$

The source terms on the right-hand side should be interpreted as the sum over all relevant matter components (e.g. photons, dark matter, baryons, etc.). The Poisson equation takes a particularly simple form if we introduce the comoving gauge density contrast

$$\nabla^2\Phi = 4\pi G a^2 \bar{\rho} \Delta . \quad (4.4.172)$$

- From the conservation of the stress-tensor, we derived the relativistic generalisations of the continuity equation and the Euler equation

$$\delta' + 3\mathcal{H} \left(\frac{\delta P}{\delta\rho} - \frac{\bar{P}}{\bar{\rho}} \right) \delta = - \left(1 + \frac{\bar{P}}{\bar{\rho}} \right) (\nabla \cdot \mathbf{v} - 3\Phi') , \quad (4.4.173)$$

$$\mathbf{v}' + 3\mathcal{H} \left(\frac{1}{3} - \frac{\bar{P}'}{\bar{\rho}'} \right) \mathbf{v} = - \frac{\nabla\delta P}{\bar{\rho} + \bar{P}} - \nabla\Phi . \quad (4.4.174)$$

Inflation: the basic picture

Key steps to understand how perturbations are generated by inflation:

- At early time all perturbation modes of interest are casually connected, i.e. correspond to k larger than the horizon: $k > aH$.
- On these (small) scales) perturbations in the inflaton field are described by a collection of harmonic oscillators
- These perturbations have quantum nature and can be followed using quantum mechanics canonical quantification. Their amplitudes have a non-zero variance:


$$\langle |\delta\phi_k|^2 \rangle \equiv \langle 0 | |\delta\phi_k|^2 | 0 \rangle$$

- Inflaton perturbations induce comoving curvature fluctuations. In the spatially flat gauge

$$\mathcal{R} = -\frac{\mathcal{H}}{\dot{\phi}} \delta\phi$$

- Thus the curvature (gauge-invariant) fluctuations have a non-zero variance:

$$\langle |\mathcal{R}_k|^2 \rangle = \left(\frac{\mathcal{H}}{\dot{\phi}} \right)^2 \langle |\delta\phi_k|^2 \rangle$$



Mukahnov-Sasaki equation

Classical inflaton field fluctuations:

Let us first see how the **inflaton field action** can be used to derive the inflaton perturbations. The action is:

$$S = \int d\tau d^3x \sqrt{-g} \left[\frac{1}{2} g^{\mu\nu} \partial_\mu \phi \partial_\nu \phi - V(\phi) \right]$$

(the integrand function is the Lagrangian density). Evaluating for a unperturbed FLRW metric one gets (prove this):

$$S = \int d\tau d^3x \left[\frac{1}{2} a^2 ((\phi')^2 - (\nabla\phi)^2) - a^4 V(\phi) \right]$$

To introduce perturbations it is convenient to write them in the following way:

$$\phi(\tau, \mathbf{x}) = \bar{\phi}(\tau) + \frac{f(\tau, \mathbf{x})}{a(\tau)}$$

To derive an equation of motion for the perturbation $f(\tau, \mathbf{x})$ one usually does:

- Use $\phi(\tau, \mathbf{x})$ in the action S .
- Expand the action up to 2nd order in the fluctuations f
- Collect all 1st order and 2nd order action terms in 2 separate actions: $S^{(1)}$ and $S^{(2)}$.
- Apply the Euler-Lagrange equations to both actions.

Mukahnov-Sasaki equation

Classical inflaton field fluctuations:

The result for using the action, $S^{(1)}$, gives the Klein-Gordon equation for the background field:

$$\bar{\phi}'' + 2\mathcal{H}\bar{\phi}' + a^2 V_{,\phi} = 0$$

From the $S^{(2)}$, which can be approximated by (see Baumann Sect. 6.2),

$$S^{(2)} \approx \int d\tau d^3x \frac{1}{2} \left[(f')^2 - (\nabla f)^2 + \frac{a''}{a} f^2 \right]$$

the Euler-Lagrange equation gives the so called **Mukahnov-Sasaki** equation

$$f'' - \nabla^2 f - \frac{a''}{a} f = 0 \quad (\text{real space-time})$$

$$f_k'' + \left(k^2 - \frac{a''}{a} \right) f_k = 0 \quad (\text{fourier space-time})$$

This has an exact solution of the form:

$$f_k(\tau) = \alpha \frac{e^{-ik\tau}}{\sqrt{2k}} \left(1 - \frac{i}{k\tau} \right) + \beta \frac{e^{ik\tau}}{\sqrt{2k}} \left(1 + \frac{i}{k\tau} \right)$$

19

Mukahnov-Sasaki equation

Classical inflaton field fluctuations:

where α , and β are set by imposing as initial conditions a plane-wave solution at early times, $\tau \rightarrow 0$. The solution, assuming a pure de Sitter space where $a = e^{Ht}$

$$\tau = \int^t e^{-Ht} dt = -H^{-1} e^{-Ht} = -\frac{1}{aH} \quad ; \quad \frac{a''}{a} = \frac{2}{\tau^2}$$

is then

$$f_k(\tau) = \frac{e^{-ik\tau}}{\sqrt{2k}} \left(1 - \frac{i}{k\tau} \right)$$

On **sub-horizon scales**, $k^2 \gg a''/a \approx 2\mathcal{H}^2$, the M-S equation becomes

$$f_k'' + k^2 f_k \approx 0$$

which is a classical harmonic oscillator with spatial frequency $\omega(k) = k$.

However we expect these fluctuations to be of quantum mechanics (QM) nature. To treat this one applies the canonical formalism of QM to the classical harmonic oscillator.

20

Quantum fluctuations in de Sitter space

Canonical quantization of the inflaton fluctuations:

One proceeds as for the harmonic oscillator theory in QM. The relevant classical quantities in the action $S^{(2)}$ are the:

- Inflaton fluctuation: $f = a\delta\phi$
- Momentum conjugate of f : $\pi \equiv \frac{\partial \mathcal{L}}{\partial f'} = f'$

One then **promotes the fields $f(\tau, \mathbf{x})$ and $\pi(\tau, \mathbf{x})$ to quantum operators** that satisfy the following commutation rules:

$$\begin{aligned} [\hat{f}(\tau, \mathbf{x}), \hat{\pi}(\tau, \mathbf{x}')] &= i\delta(\mathbf{x} - \mathbf{x}') \\ [\hat{f}_{\mathbf{k}}(\tau), \hat{\pi}_{\mathbf{k}'}(\tau)] &= \int \frac{d^3x}{(2\pi)^{3/2}} \int \frac{d^3x'}{(2\pi)^{3/2}} \underbrace{[\hat{f}(\tau, \mathbf{x}), \hat{\pi}(\tau, \mathbf{x}')] }_{i\delta(\mathbf{x} - \mathbf{x}')} e^{-i\mathbf{k}\cdot\mathbf{x}} e^{-i\mathbf{k}'\cdot\mathbf{x}'} \\ &= i \int \frac{d^3x}{(2\pi)^3} e^{-i(\mathbf{k}+\mathbf{k}')\cdot\mathbf{x}} \\ &= i\delta(\mathbf{k} + \mathbf{k}') , \end{aligned}$$

i.e. they commute in real and fourier spaces for $x \neq x'$ and $k \neq -k'$, respectively²¹

Quantum fluctuations in de Sitter space

Canonical quantization of the inflaton fluctuations:

The inflaton perturbation operator can then be written in terms of the creation and annihilation operators:

$$\hat{f}_{\mathbf{k}}(\tau) = f_{\mathbf{k}}(\tau) \hat{a}_{\mathbf{k}} + f_{\mathbf{k}}^*(\tau) \hat{a}_{\mathbf{k}}^\dagger$$

where $f_{\mathbf{k}}(\tau)$ and $f_{\mathbf{k}}^*(\tau)$ are the solution of the M-S equation,

$$f_{\mathbf{k}}'' + \omega_{\mathbf{k}}^2(\tau) f_{\mathbf{k}} = 0 , \quad \text{where} \quad \omega_{\mathbf{k}}^2(\tau) \equiv k^2 - \frac{a''}{a}$$

The creation and annihilation operators verify

$$[\hat{a}_{\mathbf{k}}, \hat{a}_{\mathbf{k}'}^\dagger] = \delta(\mathbf{k} + \mathbf{k}')$$

The quantum states (in the Hilbert space) are constructed by defining a **vacuum state** $|0\rangle$ via the condition $\hat{a}_{\mathbf{k}}|0\rangle = 0$.

Excited states of the inflaton perturbation are created using the usual creation rule:

$$|m_{\mathbf{k}_1}, n_{\mathbf{k}_2}, \dots\rangle = \frac{1}{\sqrt{m!n!\dots}} \left[(a_{\mathbf{k}_1}^\dagger)^m (a_{\mathbf{k}_2}^\dagger)^n \dots \right] |0\rangle$$

Quantum fluctuations in de Sitter space

Quantum fluctuations about the zero point (vacuum state):

Finally one can obtain inflaton perturbation operator spectrum by computing the mean and variance expectation values about the vacuum state $|0\rangle$. One has:

$$\hat{f}(\tau, \mathbf{x}) = \int \frac{d^3k}{(2\pi)^{3/2}} \left[f_k(\tau) \hat{a}_{\mathbf{k}} + f_k^*(\tau) \hat{a}_{\mathbf{k}}^\dagger \right] e^{i\mathbf{k}\cdot\mathbf{x}} .$$

The expectation value for $\langle \hat{f} \rangle = 0$ naturally, but the variance does not. One has:

$$\begin{aligned} \langle |\hat{f}|^2 \rangle &\equiv \langle 0 | \hat{f}^\dagger(\tau, \mathbf{0}) \hat{f}(\tau, \mathbf{0}) | 0 \rangle \\ &= \int \frac{d^3k}{(2\pi)^{3/2}} \int \frac{d^3k'}{(2\pi)^{3/2}} \langle 0 | (f_k^*(\tau) \hat{a}_{\mathbf{k}}^\dagger + f_k(\tau) \hat{a}_{\mathbf{k}}) (f_{k'}(\tau) \hat{a}_{\mathbf{k}'} + f_{k'}^*(\tau) \hat{a}_{\mathbf{k}'}^\dagger) | 0 \rangle \\ &= \int \frac{d^3k}{(2\pi)^{3/2}} \int \frac{d^3k'}{(2\pi)^{3/2}} f_k(\tau) f_{k'}^*(\tau) \langle 0 | [\hat{a}_{\mathbf{k}}, \hat{a}_{\mathbf{k}'}^\dagger] | 0 \rangle \\ &= \int \frac{d^3k}{(2\pi)^3} |f_k(\tau)|^2 \\ &= \int d \ln k \frac{k^3}{2\pi^2} |f_k(\tau)|^2 . \end{aligned}$$

23

Quantum fluctuations in de Sitter space

Quantum fluctuations about the zero point (vacuum state):

One defines the dimensionless power spectrum of the inflaton fluctuations as

$$\Delta_f^2(k, \tau) \equiv \frac{k^3}{2\pi^2} |f_k(\tau)|^2$$

This means that the classical solution $f_k(\tau)$ determines the variance of the quantum fluctuations. Given the relation between the fluctuation f and the inflaton field, $\delta\phi = f / a$ one has:

$$\Delta_{\delta\phi}^2(k, \tau) = a^{-2} \Delta_f^2(k, \tau) = \left(\frac{H}{2\pi} \right)^2 \left(1 + \left(\frac{k}{aH} \right)^2 \right) \xrightarrow{\text{superhorizon}} \left(\frac{H}{2\pi} \right)^2$$

So at horizon crossing one can use the following approximation:

$$\Delta_{\delta\phi}^2(k) \approx \left(\frac{H}{2\pi} \right)^2 \Big|_{k=aH}$$

Going back to the relation between the inflaton fluctuation and the curvature fluctuations,

$$\langle |\mathcal{R}_k|^2 \rangle = \left(\frac{\mathcal{H}}{\dot{\phi}'} \right)^2 \langle |\delta\phi_k|^2 \rangle$$

24

Quantum fluctuations in de Sitter space

Quantum fluctuations about the zero point (vacuum state):

The power spectra of these quantities is related via:

$$\Delta_{\mathcal{R}}^2 = \frac{1}{2\varepsilon} \frac{\Delta_{\delta\phi}^2}{M_{\text{pl}}^2}, \quad \text{where } \varepsilon = \frac{\frac{1}{2}\dot{\phi}^2}{M_{\text{pl}}^2 H^2}$$

So the power spectrum of the comoving curvature fluctuations is:

$$\Delta_{\mathcal{R}}^2(k) = \frac{1}{8\pi^2} \frac{1}{\varepsilon} \frac{H^2}{M_{\text{pl}}^2} \Big|_{k=aH}$$

which is gauge invariant and remains constant when the wavenumber k leaves the horizon scale ($k_H = aH$) during inflation.

Since the right hand side of the power spectra is evaluated at horizon crossing, $k = aH$, the power spectrum is a purely function of k . It is often useful to model this k dependence as:

$$\Delta_{\mathcal{R}}^2(k) \equiv A_s \left(\frac{k}{k_\star} \right)^{n_s - 1}$$

25

Quantum fluctuations in de Sitter space

Comoving curvature power spectrum:

The measured amplitude of the scalar spectrum at $k_\star = 0.05 \text{ Mpc}^{-1}$ is

$$A_s = (2.196 \pm 0.060) \times 10^{-9}. \quad (6.5.60)$$

To quantify the deviation from scale-invariance we have introduced the *scalar spectral index*

$$n_s - 1 \equiv \frac{d \ln \Delta_{\mathcal{R}}^2}{d \ln k}, \quad (6.5.61)$$

where the right-hand side is evaluated at $k = k_\star$ and $n_s = 1$ corresponds to perfect scale-invariance. We can split (6.5.61) into two factors

$$\frac{d \ln \Delta_{\mathcal{R}}^2}{d \ln k} = \frac{d \ln \Delta_{\mathcal{R}}^2}{dN} \times \frac{dN}{d \ln k}. \quad (6.5.62)$$

The derivative with respect to e -folds is

$$\frac{d \ln \Delta_{\mathcal{R}}^2}{dN} = 2 \frac{d \ln H}{dN} - \frac{d \ln \varepsilon}{dN}. \quad (6.5.63)$$

The first term is just -2ε and the second term is $-\eta$ (see Chapter 2). The second factor in (6.5.62) is evaluated by recalling the horizon crossing condition $k = aH$, or

$$\ln k = N + \ln H. \quad (6.5.64)$$

Hence, we have

$$\frac{dN}{d \ln k} = \left[\frac{d \ln k}{dN} \right]^{-1} = \left[1 + \frac{d \ln H}{dN} \right]^{-1} \approx 1 + \varepsilon. \quad (6.5.65)$$


To first order in the Hubble slow-roll parameters, we therefore find

$$n_s - 1 = -2\varepsilon - \eta. \quad (6.5.66)$$


26

CMB Sky statistics: The
angular power spectrum


Temperature fluctuation field



Why a ellipse-like map?



Why a ellipse-like map?



Temperature fluctuation field

- Decomposition of the temperature field on the sky:

$$\Theta(\hat{n}) = \Delta T/T(\theta, \phi) = \sum_{\ell=0}^{\infty} \sum_{m=-\ell}^{\ell} a_{\ell m} Y_{\ell m}(\theta, \phi)$$

- the $a_{\ell m}$, the decomposition coefficients, are called multi-pole moments:

$$a_{\ell m} = \int Y_{\ell m}^*(\theta', \phi') \frac{\Delta T}{T}(\theta', \phi') d\Omega'$$

these can be computed directly from the sky map. Are generally complex quantities.

Spherical harmonics


$$Y_{\ell}^m(\theta, \varphi) = \sqrt{\frac{(2\ell + 1)(\ell - m)!}{4\pi(\ell + m)!}} \cdot e^{im\varphi} \cdot P_{\ell}^m(\cos \theta)$$

$$\int_{\theta=0}^{\pi} \int_{\varphi=0}^{2\pi} Y_{\ell}^m Y_{\ell'}^{m'}* d\Omega = \delta_{\ell\ell'} \delta_{mm'} \quad d\Omega = \sin \theta d\varphi d\theta$$

Spherical harmonics:


Form a vector basis on the sphere

$\ell=1$




Made by Matthias Bartelmann

$\ell=2$




Made by Matthias Bartelmann

$\ell=3$




Made by Matthias Bartelmann

$\ell=4$




Made by Matthias Bartelmann

$l=5$




Made by Matthias Bartelmann

$l=6$



Made by Matthias Bartelmann

$\ell=7$




Made by Matthias Bartelmann

$\ell=8$

i.e. larger ℓ means shorter wavelengths


i.e. ℓ is spherical equivalent of wavenumber

$\ell \sim \pi / \theta$




Made by Matthias Bartelmann

$\ell=1$




Made by Matthias Bartelmann

$\ell=1$ plus $\ell=2$




Made by Matthias Bartelmann

$\ell=1$ plus $\ell=2$ plus $\ell=3$




Made by Matthias Bartelmann

Sum $\ell=1$ to 4




Made by Matthias Bartelmann

Sum $\ell=1$ to 5




Made by Matthias Bartelmann

Sum $\ell=1$ to 6




Made by Matthias Bartelmann

Sum $\ell=1$ to 7




Made by Matthias Bartelmann

Sum $\ell=1$ to 8




Made by Matthias Bartelmann

Sum up to some high l



Made by Matthias Bartelmann

Map with all multipoles



Angular correlation function

- the temperature fluctuation field is assumed as Gaussian Random variable. It's angular correlation function

$$C(\hat{n}, \hat{n}') \equiv \left\langle \frac{\Delta T}{T}(\hat{n}) \frac{\Delta T}{T}(\hat{n}') \right\rangle = \sum_{\ell \ell'} \sum_{m m'} \langle a_{\ell m}^* a_{\ell' m'} \rangle Y_{\ell m}^*(\hat{n}) Y_{\ell' m'}(\hat{n}')$$

fully characterizes the temperature fluctuation field (brackets denote averages over an ensemble of Universes). It is conventional to write (the $a_{\ell m}$ are not correlated):

$$\langle a_{\ell m}^* a_{\ell' m'} \rangle = C_\ell \delta_{\ell \ell'} \delta_{m m'} \quad , \quad C_\ell \equiv \langle |a_{\ell m}|^2 \rangle$$

C_ℓ is the angular power spectrum. Then we have

$$C(\hat{n}, \hat{n}') = \sum_{\ell} \frac{(2\ell + 1)}{4\pi} C_\ell P_\ell(\cos \vartheta) = C(\cos \vartheta)$$

CMB angular power spectra

- temperature fluctuation spectrum:

$$\langle a_{\ell m}^* a_{\ell' m'} \rangle = C_\ell \delta_{\ell \ell'} \delta_{m m'} \quad , \quad C_\ell \equiv \langle |a_{\ell m}|^2 \rangle$$

- Polarization and cross correlation power spectra:

$$\langle E_{\ell m}^* E_{\ell' m'} \rangle = \delta_{\ell \ell'} \delta_{m m'} C_\ell^{EE} ,$$


$$\langle B_{\ell m}^* B_{\ell' m'} \rangle = \delta_{\ell \ell'} \delta_{m m'} C_\ell^{BB} ,$$

$$\langle \Theta_{\ell m}^* E_{\ell' m'} \rangle = \delta_{\ell \ell'} \delta_{m m'} C_\ell^{\Theta E} .$$


these quantities are highly sensitive to the cosmological parameters. They can be computed theoretically and measured from sky maps. Powerful tool to constrain cosmological parameters

CMB angular power spectra

Planck



CMB angular power spectra




Sachs–Wolfe effect

$$\Delta v/v \sim \Delta T/T \sim \Phi/c^2$$

Additional effect of time dilation while potential evolves (White & Hu 1997):

$$\frac{\Delta T}{T} \sim \frac{1}{3} \frac{\Delta \Phi}{c^2}$$



The temperature fluctuations due to the so-called Sachs–Wolfe effect are due to two competing effects: (1) the redshift experienced by the photon as it climbs out of the potential well toward us and (2) the delay in the release of the radiation, leading to less cosmological redshift compared to the average CMB radiation.

The first contribution leads to a redshift of the order of:

$$\frac{\delta T_1}{T} = \frac{\delta \Phi}{c^2}$$

Sachs–Wolfe effect

The second contribution is more tricky. Because of general relativity, the proper time goes slower inside the potential well than outside. The cooling of the gas in this potential well thus also goes slower, and it therefore reaches 3000 K at a later time relative to the average Universe.

The time delay (in terms of global time t) is:

$$\frac{\delta t}{t} = -\frac{\delta \Phi}{c^2} \quad (8.7)$$

This means that 3000 K is reached at a slightly larger (global) scale parameter $a + \delta a > a$. Since in the Einstein-de-Sitter Universe we have $a \propto t^{2/3}$ we can write

$$\frac{\delta a}{a} = \frac{2}{3} \frac{\delta t}{t} = -\frac{2}{3} \frac{\delta \Phi}{c^2} \quad (8.8)$$

Now, from that point $a = (a_{\text{cmb}} + \delta a)$ until today $a = 1$ the redshift due to expansion is less by:

$$\frac{\delta z}{z} = -\frac{\delta a}{a} \quad (8.9)$$

which leads to a positive contribution to the temperature fluctuation δT that we observe today:

$$\frac{\delta T_2}{T} = -\frac{\delta z}{z} = \frac{\delta a}{a} = -\frac{2}{3} \frac{\delta \Phi}{c^2} \quad (8.10)$$

The total is the sum of both contributions:

$$\frac{\delta T}{T} = \frac{\delta T_1}{T} + \frac{\delta T_2}{T} = \frac{1}{3} \frac{\delta \Phi}{c^2} \quad (8.11)$$

Sachs–Wolfe effect

For power-law index of primary density perturbations ($n_s=1$, Harrison–Zel’dovich spectrum), the Sachs–Wolfe effect produces a flat power spectrum: $C_\ell^{\text{SW}} \sim 1/\ell(\ell+1)$

$$\begin{aligned} C_\ell &= \frac{1}{25} \int \frac{d^3k}{k^3} \mathcal{P}_{\mathcal{R}}(k) j_\ell(kx)^2 \\ &= \frac{4\pi}{25} \int_0^\infty \frac{dk}{k} \mathcal{P}_{\mathcal{R}}(k) j_\ell(kx)^2, \end{aligned} \quad (67)$$

the final result for an arbitrary primordial power spectrum $\mathcal{P}_{\mathcal{R}}(k)$.

The integral can be done for a power-law power spectrum, $\mathcal{P}(k) = A^2 k^{n-1}$. In particular, for a scale-invariant ($n = 1$) primordial power spectrum,

$$\mathcal{P}_{\mathcal{R}}(k) = \text{const.} = A^2, \quad (68)$$

we have

$$C_\ell = A^2 \frac{4\pi}{25} \int_0^\infty \frac{dk}{k} j_\ell(kx)^2 = \frac{A^2}{25} \frac{2\pi}{\ell(\ell+1)}, \quad (69)$$

since

$$\int_0^\infty \frac{dk}{k} j_\ell(kx)^2 = \frac{1}{2\ell(\ell+1)}. \quad (70)$$

We can write this as


$$\frac{\ell(\ell+1)}{2\pi} C_\ell = \frac{A^2}{25} = \text{const. (independent of } \ell) \quad (71)$$

Acoustic oscillations

- Baryons fall into dark matter potential wells: [Photon baryon fluid heats up](#)
- Radiation pressure from photons resists collapse, overcomes gravity, expands: [Photon–baryon fluid cools down](#)
- Oscillating cycles on all scales. Sound waves stop oscillating at recombination when photons and baryons decouple.

Credit: Wayne Hu

Springs:
photon
pressure



Balls:
baryon
mass

Acoustic peaks

Oscillations took place on all scales. We see temperature features from modes which had reached the extrema


- Maximally compressed regions were hotter than the average
Recombination happened later, corresponding photons experience less red-shifting by Hubble expansion: **HOT SPOT**
- Maximally rarified regions were cooler than the average
Recombination happened earlier, corresponding photons experience more red-shifting by Hubble expansion: **COLD SPOT**

Harmonic sequence, like waves in pipes or strings:



2nd harmonic: mode compresses and rarifies by recombination

3rd harmonic: mode compresses, rarifies, compresses


➔ 2nd, 3rd, .. peaks



Harmonic sequence



Credit: Wayne Hu



Modes with half the wavelengths oscillate twice as fast ($v = c/\lambda$).

➔ Peaks are equally spaced in l


Doppler shifts

Times in between maximum compression/rarefaction, modes reached maximum velocity

This produced temperature enhancements via the Doppler effect
(non-zero velocity along the line of sight)

This contributes power in between the peaks



➔ Power spectrum does not go to zero



Damping and diffusion

- Photon diffusion (Silk damping) suppresses fluctuations in the baryon-photon plasma
- Recombination does not happen instantaneously and photons execute a random walk during it. Perturbations with wavelengths which are shorter than the photon mean free path are damped (the hot and cold parts mix up)


This is same as a low-resolution instrument blurs all the details!



Thickness of the LSS is comparable to the oscillation scales

➔ Power falls off

Online C_l calculators




CMB Toolbox: <http://lambda.gsfc.nasa.gov/toolbox/>

CAMB website: <http://camb.info/>

CMBFast website: <http://www.cmbfast.org/>

temperature power spectrum: parameter dependence



There are model degeneracies among parameters.

Exercise:

Go online to <http://lambda.gsfc.nasa.gov/toolbox/> and use the CAMB online tool to assess the effect of the following parameters on the temperature angular power spectrum of the CMB; $\Omega_b h^2$; $\Omega_m h^2$, Ω_Λ .

Reprinted from: Lecture Notes on CMB Theory: From Nucleosynthesis to Recombination, Wayne Hu, arXiv:0802.3688




Fig. 15. Baryons and matter. Baryons change the relative heights of the even and odd peaks through their inertia in the plasma. The matter-radiation ratio also changes the overall amplitude of the oscillations from driving effects. Adapted from [Hu and Dodelson \(2002\)](#).

Exercise:

Go online to <http://lambda.gsfc.nasa.gov/toolbox/> and use the CAMB online tool to assess the effect of the following parameters on the temperature angular power spectrum of the CMB; $\Omega_b h^2$; $\Omega_m h^2$, Ω_Λ .

Reprinted from: Lecture Notes on CMB Theory: From Nucleosynthesis to Recombination, Wayne Hu, arXiv:0802.3688




Fig. 14. Curvature and dark energy. Given a fixed physical scale for the acoustic peaks (fixed $\Omega_b h^2$ and $\Omega_m h^2$) the observed angular position of the peaks provides a measure of the angular diameter distance and the parameters it depends on: curvature, dark energy density and dark energy equation of state. Changes at low ℓ multipoles are due to the decay of the gravitational potential after matter domination from the integrated Sachs-Wolfe effect.

CMB parameter cheat sheet

