



# Toxicity of the insecticides spinosad and indoxacarb to the non-target aquatic midge *Chironomus riparius*

Hugo R. Monteiro<sup>a,b,c,\*</sup>, João L.T. Pestana<sup>a</sup>, Sara C. Novais<sup>b</sup>, Amadeu M.V.M. Soares<sup>a</sup>, Marco F.L. Lemos<sup>b</sup>

<sup>a</sup> Department of Biology & CESAM, University of Aveiro, Aveiro, Portugal


<sup>b</sup> MARE – Marine and Environmental Sciences Centre, ESTM, Instituto Politécnico de Leiria, Peniche, Portugal

<sup>c</sup> Department of Biochemistry and Microbiology, Laboratory for Microbiology, Ghent University, Ghent, Belgium

## HIGHLIGHTS

- Responses of *C. riparius* to spinosad and indoxacarb exposure were investigated.
- Exposure to both insecticides compromised *C. riparius* life-history traits.
- Both insecticides induced defense mechanisms and cellular oxygen consumption.
- Non-target aquatic insect species may be impacted by spinosad and indoxacarb use.

## GRAPHICAL ABSTRACT



## ARTICLE INFO

### Article history:

Received 30 November 2018

Received in revised form 18 February 2019

Accepted 19 February 2019

Available online 20 February 2019

Editor: Damian Barcelo

### Keywords:

Aquatic invertebrates  
Biochemical biomarkers  
Insecticides  
Life-history effects  
Neurotoxicity

## ABSTRACT

Spinosad and indoxacarb are two relatively new insecticides mainly used in agriculture to control insect pests. However, at their current application rates, non-target aquatic insect species may also be impacted. In this study, larvae of the non-biting midge *Chironomus riparius* were exposed in the laboratory to both insecticides and their effects evaluated at the organismal level, using standard ecotoxicological tests, and at the biochemical level, by monitoring specific oxidative stress, neuronal, and energy metabolism biomarkers. Chronic exposure to both insecticides compromised growth and emergence of *C. riparius*. Short-term exposures revealed alterations at biochemical level that might be related to the toxicological targets of both insecticides. Growth and development time were the most sensitive endpoints at individual level for both pesticides, while at the biochemical level, the electron transport system activity was the most sensitive biomarker for spinosad exposure, suggesting an increase in energy demands associated with the activation of defense mechanisms. Glutathione-S-transferase was the most sensitive biomarker for indoxacarb exposure, underlining the role of this enzyme in the detoxification of indoxacarb. Additionally, changes in lactate dehydrogenase and glutathione peroxidase activities were observed for both insecticides, and evidences of oxidative damage were found for spinosad. This study contributes to the growing knowledge on sublethal effects of novel insecticides on non-target aquatic invertebrates and strengthens the usefulness of biochemical biomarkers to support the interpretation of their potentially deleterious effects on aquatic insects near agricultural fields.

© 2019 Elsevier B.V. All rights reserved.

\* Corresponding author at: Department of Biology & CESAM, University of Aveiro, Aveiro, Portugal.  
E-mail address: [hugoricardo@ua.pt](mailto:hugoricardo@ua.pt) (H.R. Monteiro).

## 1. Introduction

One of the goals in integrated pest management is to find tailor-made and effective pesticides for specific pests while keeping adverse consequences on non-target species to a minimum (Chitgar and Ghadamyari, 2012; Stara et al., 2010; Wilkinson et al., 1979). Some non-target aquatic insects play vital roles in freshwater ecosystems, and are regularly subjected to pesticide exposure through runoff, drift, or leaching from adjacent agricultural fields (Cerejeira et al., 2003; Schulz, 2004). Ecotoxicological effects of pesticides seen on higher levels of organization are often preceded by quantifiable alterations at biochemical levels, and assessing earlier sub-organismal endpoints on key species may provide insights on the long-term consequences for natural populations (Lemos et al., 2010), hopefully providing regulators with early-warning tools for risk assessment.

Spinosad and indoxacarb are neurotoxic insecticides with distinct modes of action registered for agricultural use in Europe (European Commission, 2006, 2008a). However, the toxicity data for these relatively novel insecticides on aquatic invertebrates is still very limited, considering that according to regulation (EC) No 1272/2008 (European Commission, 2008b) they are both classified as very toxic to aquatic life with long lasting effects.

Spinosad targets a unique site in nicotinic acetylcholine receptors (Copping and Menn, 2000; Orr et al., 2009), causing hyperexcitation of the nervous system (Salgado, 1998; Salgado et al., 1998; Salgado and Sparks, 2005). It is very effective against several insect species (Hertlein et al., 2010), including chironomids (Bond et al., 2004; Lawler and Dritz, 2013; Pérez et al., 2007; Stevens et al., 2005). Recently, the European Food Safety Authority predicted an environmental concentration of  $26.28 \mu\text{g L}^{-1}$  on surface waters resulting from the applications on leafy and fruity vegetables (worst case scenario) (EFSA et al., 2018a). Moreover, spinosad sorbs to the sediment where it seems to be more persistent (Cleveland et al., 2002), and where many sediment-dwelling organisms, including chironomid larvae, may be affected.

Indoxacarb acts by blocking voltage-dependent sodium channels, causing nervous system shutdown (Lapied et al., 2001; Wing et al., 1998, 2000). It is effective against several insect species (Anikwe et al., 2014; Dryden et al., 2013; Oxborough et al., 2015; Pridgeon et al., 2009), but particularly to lepidopterans (Dias, 2006; Wing et al., 1998, 2000). In 2003, indoxacarb estimated environmental long-term average concentrations in surface waters were of  $3.7 \mu\text{g L}^{-1}$ , with peak values of  $13.7 \mu\text{g L}^{-1}$  (EPA, 2003). More recently, levels up to  $7.763 \mu\text{g L}^{-1}$ , resulting from indoxacarb's application in lettuce crops, were predicted for surface waters (EFSA et al., 2018b). Additionally, indoxacarb also has a relatively high log Kow of 4.65 (Dias, 2006) suggesting it has a high tendency to sorb to sediments.

The freshwater midge *Chironomus riparius* Meigen (Diptera: Chironomidae) is a widely used model organism in ecotoxicology testing (Weltje et al., 2010) mainly due to its ecological relevance and easiness to handle in the laboratory. Additionally, *C. riparius* larvae have been previously used as a model to evaluate biochemical responses of insecticide exposure (Monteiro et al., 2019; Rodrigues et al., 2015a, 2015b).

The main aim of this study was to investigate organismal and biochemical effects of two neurotoxic insecticides on the aquatic invertebrate *C. riparius*. For that, environmentally relevant concentrations of spinosad and indoxacarb were used to investigate sub-lethal effects in terms of larval growth, development, and emergence using standard ecotoxicological tests. Moreover, the effects of these insecticides at the biochemical level were determined to evaluate their possible relation to the effects observed at the organismal level and their potential use in biomonitoring studies.

Biochemical biomarkers associated with key physiological functions were selected: activity of the antioxidant enzymes superoxide dismutase (SOD), catalase (CAT), glutathione peroxidase (GPx), and glutathione reductase (GR), essential in the protection against reactive oxygen

species (ROS) induced by the exposure to xenobiotics; DNA damage and lipid peroxidation (LPO), as indicators of oxidative damage, resulting from excessive ROS; activity of glutathione-S-transferase (GST), a phase II biotransformation enzyme, involved in the detoxification of xenobiotics; acetylcholinesterase (AChE) activity, related to the cholinergic neurotransmission, as a biomarker of neuromuscular toxicity; the activities of lactate dehydrogenase (LDH) and electron transport system (ETS) were assessed as measures of energy metabolism.

## 2. Material and methods

### 2.1. Test organism

*Chironomus riparius* larvae were collected from a laboratory culture long established at the University of Aveiro, Portugal. Larvae are kept in plastic aquaria filled with a fine layer of washed and burnt river sand (<1 mm) and American Society for Testing Materials (ASTM) hard water. Cultures were maintained at 20 °C with a photoperiod of 16:8 h light-dark, with a constant inflow of air. Larvae are fed *ad libitum* with macerated fish food, Tetramin® (Melle, Germany).

### 2.2. Test chemicals

Spinosad (CAS number 168316-95-8), a mixture of spinosyns A and D (Crouse et al., 2001), two byproducts of the fermentation of *Saccharopolyspora spinosa* (Actinomycetales: Pseudonocardiales) (Mertz and Yao, 1990; Thompson et al., 2000), and indoxacarb (CAS number 144171-61-9), an oxadiazine insecticide, were acquired from Sigma-Aldrich, UK. Stock solutions of the pesticides were prepared in ethanol (spinosad) and acetone (indoxacarb). Experimental solutions were prepared by diluting the stock solutions in ASTM hard water, and the final solvent concentration was kept at 0.01%.

### 2.3. Acute toxicity tests

Acute lethal toxicity was assessed following OECD guideline 235 (OECD, 2011) with water only exposures in crystalizing dishes, using 1st instar larvae. Larvae were exposed to concentrations of spinosad of 0 (solvent control), 1, 2, 4, 8, 16, 32, 64, 128, and  $256 \mu\text{g L}^{-1}$ , and to 0, 0.5, 1, 2, 4, 8, 16, 32, 64, and  $128 \mu\text{g L}^{-1}$  of indoxacarb. After 48 h of exposure, mortality was checked by mechanical stimulation. To halt possible photodegradation of the chemicals, crystalizing dishes were protected from the light during the test. The test was executed at  $20 \pm 1$  °C, and larvae were not fed during the exposure.

### 2.4. Chronic toxicity tests

A 28-day chronic test was performed according to the OECD guideline 219 (OECD, 2004). First instar larvae of *C. riparius* (2 days old) were exposed to 0 (negative and solvent control), 0.5, 1.28, 3.2, 8, and  $20 \mu\text{g L}^{-1}$  of spinosad in 150 mL of medium and layer of 1.5 cm of sediment (commercial river sand washed, sieved, and burnt) in 200 mL glass vessels. A similar setup was made for indoxacarb, using 0, 1, 2, 4, and  $8 \mu\text{g L}^{-1}$  treatments. Five larvae were used in each replicate, and five replicates were used for larval growth determination, while eight replicates were used for emergence endpoints. After ten days of exposure larval growth was determined by measuring body length of the larvae with the aid of a stereomicroscope fitted with a calibrated micrometer and growth was calculated by subtracting the mean body length at the beginning (pool of 25 larvae of initial size). In the eight remaining replicates, adult *C. riparius* were collected daily, their gender determined and stored in 70% ethanol. Afterwards, adult midges were dried at 50 °C for 24 h and weighed with a microbalance (Mettler UMT2).

The tests were performed under the same conditions described for culturing:  $20 \pm 1$  °C with 16:8 h light:dark cycle with gentle aeration.

Organisms were fed every two days at a ration of 0.5 mg Tetramin® larvae<sup>-1</sup> day<sup>-1</sup>, and physicochemical parameters were checked throughout the experiment.

## 2.5. Biomarkers

For the determination of the biochemical biomarkers, tests were initiated with 8 days old larvae (3rd instar). The concentrations used in these bioassays were 0, 0.5, 2, and 8 µg L<sup>-1</sup> for spinosad, and 0, 2, 4, and 8 µg L<sup>-1</sup> for indoxacarb. Each crystalizing dish contained ten larvae, 80 mL of experimental solution, and a fine layer of sediment. After 48 h, organisms from two replicates of the same treatment were pooled to give a total of twenty organisms per replicate. Six pooled replicates were used per treatment for spinosad, and seven pooled replicates for indoxacarb. After exposure, larvae were collected, excess water was gently removed with a filter paper, and organisms weighed, frozen with liquid nitrogen, and stored at -80 °C until further processing.

Samples were subsequently homogenized in 800 µL of 0.1 M of K-phosphate buffer (pH = 7.4) using an Ystral d-79282 homogenizer. This homogenate was divided into aliquots for ETS, LPO, and DNA damage determination. To LPO aliquot, 4% 2,6-Di-tert-butyl-4-methylphenol (BHT) in methanol was added to prevent subsequent lipid oxidation of the samples (Aloísio Torres et al., 2002). These three aliquots were immediately stored at -80 °C until used. The remaining homogenate was centrifuged at 10,000g for 20 min at 4 °C and the supernatant (post-mitochondrial supernatant) was collected and divided into aliquots for SOD, CAT, GST, GR, GPx, AChE, and LDH activities determination and for protein quantification. In every assay, reaction blanks were performed using K-phosphate buffer instead of the sample and all spectrophotometric measurements were made at 25 °C using a Synergy H1 Hybrid Multi-Mode microplate reader (BioTek® Instruments, Vermont, USA).

### 2.5.1. Protein quantification

Protein concentration was assessed following the Bradford protocol adapted to microplate, using γ-globuline as standard. Prior to AChE, CAT, GR, GPx, GST, and LDH activities determination, protein concentration was adjusted to approximately 0.8 mg L<sup>-1</sup>. For these biomarkers, the exact protein concentration of the dilution was measured again at the end of the experiment.

### 2.5.2. Detoxification, oxidative stress and oxidative damage biomarkers

SOD activity was determined by following the method described by McCord and Fridovich (1969) adapted to microplate. Cytochrome c reduction was followed for 5 min at 550 nm, and results are expressed as SOD units (U) mg<sup>-1</sup> protein. The determination of CAT activity was made according to Clairborne (1985). The consumption of H<sub>2</sub>O<sub>2</sub> was assessed at 240 nm for 1 min, and results are expressed in µmol min<sup>-1</sup> mg<sup>-1</sup> of protein. For the assessment of GR activity, the method described by Cribb et al. (1989) was used. The oxidation of NADPH was monitored at 340 nm during 1 min, and results are expressed in nmol min<sup>-1</sup> mg<sup>-1</sup> of protein. Regarding GPx activity, it was determined by monitoring the oxidation of NADPH at 340 nm for 3 min, as a result of GR conversion of GSSG to GSH (Mohandas et al., 1984). Results are expressed in nmol min<sup>-1</sup> mg<sup>-1</sup> of protein. An adaption of Habig et al. (1974) protocol to microplate was used to determine GST activity. The formation of glutathione dinitrobenzene was measured at 340 nm during 3 min, and results are expressed in nmol min<sup>-1</sup> mg<sup>-1</sup> of protein. LPO levels were measured using thiobarbituric acid reactive substances (TBARS) assay (Bird and Draper, 1984; Ohkawa et al., 1979). Absorbance was read at 535 nm and results are expressed in nmol TBARS g<sup>-1</sup> of wet weight. DNA damage was assessed following the protocols described by de Lafontaine et al. (2000) and Olive (1988). Fluorescence was measured using an excitation/emission wavelength of 360/460 nm, and results are expressed as ng of damaged DNA mg<sup>-1</sup> of wet weight.

### 2.5.3. Neurotransmission and energy related biomarkers

Effects of spinosad and indoxacarb on cholinergic neurotransmission were evaluated monitoring AChE activity, following Ellman's method (Ellman et al., 1961) adapted to microplate (Guilhermino et al., 1996). The absorbance was read at 414 nm for 5 min, and results are expressed in nmol min<sup>-1</sup> mg<sup>-1</sup> of protein. To determine the activity of anaerobic metabolism-related enzyme LDH, oxidation of NADH was monitored at 340 nm as proposed by Vassault (1983) and Diamantino et al. (2001). Results are expressed in nmol min<sup>-1</sup> mg<sup>-1</sup> of protein. ETS activity was determined following De Coen and Janssen (1997) with some adaptations (Rodrigues et al., 2015b). Absorbance was read at 490 nm for 5 min, and results are expressed mJ h<sup>-1</sup> mg of protein<sup>-1</sup>.

## 2.6. Statistical analysis

Effects of insecticide exposure on life history and biochemical endpoints were evaluated by one-way analysis of variance (ANOVA) followed by a Dunnett's post hoc test to determine statistically significant differences between solvent controls and treatments, and/or by a test for linear trend to discriminate if there is a linear increase or decrease in response as the concentration increases. Data were checked for residual normality using D'Agostino-Pearson and Shapiro-Wilk normality tests and for homoscedasticity with Brown-Forsythe test. Unpaired *t*-tests did not find differences between negative and solvent controls for any of the endpoints analysed, therefore solvent control was used as the control for all analysis. Spinosad's DNA damage data were log-transformed to correct for normality. For spinosad LPO data and for indoxacarb percentage of emergence data, transformations did not correct for normality, but since homogeneity of variances was verified, one-way ANOVA was executed. Since all larvae in the spinosad chronic test exposed to 20 µg L<sup>-1</sup> died, this treatment was excluded from analysis. Statistical analysis was made in GraphPad Prism® 7 for Mac and significance level was set at 0.05.

## 3. Results

### 3.1. Spinosad

For spinosad, in the highest concentration tested, there was 40% mortality after 48 h of exposure (Supplementary data, Table I). Because of the gradient of concentrations used for spinosad, the 48 h LC50 could not be estimated and thus is higher than 256 µg L<sup>-1</sup>. Concerning the chronic bioassay, at day 10 no larvae were alive at the highest concentration tested (20 µg L<sup>-1</sup>) while 92% of the larvae were recovered from the control. Additionally, at day 10 of exposure, statically significant differences were found for growth between control and the 8 µg L<sup>-1</sup> treatment ( $F_{(4,20)} = 7.640, p < 0.001$ ) (Table I). Regarding emergence parameters, there was a significant increase in time to emergence at 8 µg L<sup>-1</sup> for both males ( $F_{(4,27)} = 3.831, p < 0.05$ ) and females ( $F_{(4,26)} = 3.606, p < 0.05$ ) (Table I). No adults have emerged in the 20 µg L<sup>-1</sup> treatment, and although overall ANOVA was not significant for the remaining treatments ( $F_{(4,34)} = 2.295, p = 0.079$ ), Dunnett's test discriminated differences between control and 8 µg L<sup>-1</sup> treatments in terms of percentage of emerged adults. No effects were found on adult weight (NOEC = 8 µg L<sup>-1</sup>) (Table II).

To what concerns biochemical biomarkers, exposure to spinosad significantly increased GPx activity in *C. riparius* larvae in the highest concentration ( $F_{(3,20)} = 7.601, p < 0.01$ ; Fig. 2b), and LDH activity in the 2 µg L<sup>-1</sup> treatment ( $F_{(3,19)} = 8.357, p = 0.001$ ; Fig. 1d). There was also a significant increase in LPO at the two highest concentrations tested ( $F_{(3,20)} = 4.87, p < 0.05$ ; Fig. 1a) and, although not significant, DNA damage also increased in the same treatments ( $F_{(3,20)} = 2.651, p = 0.077$ , Fig. 1b). ETS activity was the most sensitive biomarker, with a significant increase observed for all tested concentration ( $F_{(3,20)} = 31.76, p < 0.001$ ; LOEC = 0.5 µg L<sup>-1</sup>, Fig. 1e). Moreover, this increase was concentration-dependent ( $r^2 = 0.83, p < 0.001$ ). No significant alterations were detected for AChE, CAT, GR, GST, and SOD activities (Figs. 1–2).

**Table I**  
Growth ( $n = 5$ ) and emergence endpoints ( $n = 8$ ) of *Chironomus riparius* larvae exposed to spinosad. All values are presented as mean  $\pm$  SEM. An asterisk denotes statistically significant differences to the control treatment ( $0 \mu\text{g L}^{-1}$ ;  $p < 0.05$ , ANOVA, Dunnett's test). A number sign denotes statistically significant differences to the control treatment ( $0 \mu\text{g L}^{-1}$ ; Dunnett's test) when overall ANOVA is not significant ( $p = 0.079$ ).

Spinosad concentrations ( $\mu\text{g L}^{-1}$ )	Growth (mm)	Total emergents (%)	Development time (days)	
			Males	Females
0	12.3 $\pm$ 0.2	80.0 $\pm$ 7.6	15.3 $\pm$ 0.4	16.8 $\pm$ 0.4
0.5	11.6 $\pm$ 0.3	65.0 $\pm$ 9.1	15.2 $\pm$ 0.3	17.1 $\pm$ 0.4
1.28	11.7 $\pm$ 0.3	65.0 $\pm$ 7.3	15.3 $\pm$ 0.3	17.2 $\pm$ 0.3
3.2	10.9 $\pm$ 0.5	72.5 $\pm$ 8.4	15.0 $\pm$ 0.3	17.9 $\pm$ 0.6
8	8.8 $\pm$ 0.9*	45.0 $\pm$ 9.8#	17.1 $\pm$ 0.7*	19.9 $\pm$ 1.1*
20	N.C.	N.C.	N.C.	N.C.

N.C. – not calculated due to 100% mortality.

**Table II**  
Adult weight of *Chironomus riparius* exposed as larvae to spinosad. All values are presented as mean  $\pm$  SEM,  $n = 8$ .

Spinosad concentrations ( $\mu\text{g L}^{-1}$ )	Males dry weight (mg)	Females dry weight (mg)
0	0.542 $\pm$ 0.012	1.084 $\pm$ 0.037
0.5	0.540 $\pm$ 0.013	1.084 $\pm$ 0.019
1.28	0.568 $\pm$ 0.017	1.063 $\pm$ 0.036
3.2	0.524 $\pm$ 0.014	1.035 $\pm$ 0.033
8	0.521 $\pm$ 0.027	1.118 $\pm$ 0.091
20	N.C.	N.C.

N.C. – not calculated due to 100% mortality.

### 3.2. Indoxacarb


In acute tests and for the highest concentration of indoxacarb tested, there was 47% mortality after 48 h of exposure (Supplementary data, Table I). The 48 h LC50 of indoxacarb was estimated to be higher than  $128 \mu\text{g L}^{-1}$ . Chronic exposure to indoxacarb led to a decrease in larval growth in the highest concentration tested ( $8 \mu\text{g L}^{-1}$ ;  $F_{(4,19)} = 4.746$ ,

$p < 0.01$ ) (Table III). Moreover, indoxacarb exposure led to a delay in emergence of males ( $F_{(4,32)} = 11.96$ ,  $p < 0.001$ ) and females ( $F_{(4,33)} = 6.031$ ,  $p < 0.001$ ). No effects were observed for the percentage of emerged adults (Table III) nor for adult weight (NOEC =  $8 \mu\text{g L}^{-1}$ ) (Table IV).


Regarding biochemical biomarkers, GPx activity increased in the highest concentration tested ( $F_{(3,24)} = 5.055$ ,  $p < 0.01$ ; Fig. 4b). Exposure to indoxacarb significantly increased LDH activity in the highest concentration tested ( $F_{(3,23)} = 3.331$ ,  $p < 0.05$ ; Fig. 3d), and this increase was dose-dependent ( $r^2 = 0.30$ ,  $p < 0.01$ ; Fig. 3d). GST activity increased from concentration  $4 \mu\text{g L}^{-1}$  onwards ( $F_{(3,24)} = 4.81$ ,  $p < 0.01$ ; Fig. 3c). For SOD activity, although ANOVA was significant, the post hoc test did not find any significant differences between the control and the experimental treatments ( $F_{(3,24)} = 4.21$ ,  $p < 0.05$ ; Fig. 4d). No significant alterations were detected for the remaining biomarkers studied.

### 4. Discussion

Spinosad and indoxacarb are neurotoxic insecticides highly effective in controlling insect pests which were initially deemed as relatively safe for non-target species (Bacci et al., 2016; Boucher and Ashley, 1999;



**Fig. 1.** Oxidative damage, biotransformation, energetic metabolism and neuronal biomarkers in *Chironomus riparius* larvae after 48 h exposure to spinosad: a) Lipid Peroxidation; b) DNA Damage; c) Glutathione-S-Transferase; d) Lactate Dehydrogenase; e) Electron Transport System; f) Acetylcholinesterase. All values are presented as mean  $\pm$  SEM,  $n = 7$ . An asterisk denotes statistically significant differences to the control treatment ( $0 \mu\text{g L}^{-1}$ ;  $p < 0.05$ , ANOVA, Dunnett's test).



**Fig. 2.** Oxidative stress biomarkers in *Chironomus riparius* larvae after 48 h exposure to spinosad: a) Catalase; b) Glutathione Peroxidase; c) Glutathione Reductase; d) Superoxide Dismutase. All values are presented as mean  $\pm$  SEM,  $n = 7$ . An asterisk denotes statistically significant differences to the control treatment ( $0 \mu\text{g L}^{-1}$ ;  $p < 0.05$ , ANOVA, Dunnett's test).

**Table III**

Growth ( $n = 5$ ) and emergence endpoints ( $n = 8$ ) of *Chironomus riparius* larvae exposed to indoxacarb. All values are presented as mean  $\pm$  SEM. An asterisk denotes statistically significant differences to the control treatment ( $0 \mu\text{g L}^{-1}$ ;  $p < 0.05$ , ANOVA, Dunnett's test).

Indoxacarb concentrations ( $\mu\text{g L}^{-1}$ )	Growth (mm)	Total emergents (%)	Development time (days)	
			Males	Females
0	11.7 $\pm$ 0.3	90.0 $\pm$ 5.4	17.0 $\pm$ 0.3	18.4 $\pm$ 0.6
1	11.3 $\pm$ 0.1	90.0 $\pm$ 3.8	16.0 $\pm$ 0.3	19.5 $\pm$ 0.5
2	11.6 $\pm$ 0.6	85.0 $\pm$ 6.3	16.2 $\pm$ 0.3	18.5 $\pm$ 0.4
4	10.9 $\pm$ 0.3	82.5 $\pm$ 7.0	17.4 $\pm$ 0.4	19.9 $\pm$ 0.6
8	9.9 $\pm$ 0.4*	95.0 $\pm$ 3.2	19.0 $\pm$ 0.4*	22.1 $\pm$ 0.9*

Jones et al., 2005; Lahm et al., 2000; Liu and Zhang, 2012; Sarfraz et al., 2005). Although there is no recent literature available on measured levels of both chemicals in natural freshwater environments, the concentrations used in this study are within the estimated environmental levels, and clearly impaired *C. riparius* life-history traits with alterations at the biochemical level also observed.


Chronic exposures to indoxacarb and spinosad produced comparable outcomes in terms of *C. riparius* life history traits: growth reduction observed after 10 days of exposure at  $8 \mu\text{g L}^{-1}$  of spinosad and at  $8 \mu\text{g L}^{-1}$

of indoxacarb translated into a delay in development of both males and females, but interestingly, did not result in a reduction of imagoes weight. This suggests that *C. riparius* larvae were capable of recovering and reaching the desired weight, at the expense of longer development time. This trade-off is not unusual as body weight is associated with the reproductive output of chironomids (Sibley et al., 2001) - nonetheless, a delay in development time is still an important ecological driver as it can have direct consequences on synchrony of emergence and mating success, thus affecting population dynamics (Sibley et al., 1997). This is particularly relevant on protandrous species such as *C. riparius*. Still, a full life-cycle test contemplating reproductive endpoints such as fecundity and fertility could provide a better estimate of population-level effects and should be considered in future studies. The main dissimilarity observed between the effects of the two compounds at the organismal level, was that spinosad exposure also affected *C. riparius* survival: there was a reduction in the number of emerged adults at  $8 \mu\text{g L}^{-1}$ , and at  $20 \mu\text{g L}^{-1}$  no imagoes have emerged. Previous data indicated a NOEC (no observed effect concentration) of  $0.62 \mu\text{g L}^{-1}$  for *C. riparius* (EFSA et al., 2018a), however, in the present study a NOEC of  $3.2 \mu\text{g L}^{-1}$  and a LOEC (lowest observed effect concentration)

**Table IV**

Adult weight of *Chironomus riparius* exposed as larvae to indoxacarb. All values are presented as mean  $\pm$  SEM,  $n = 8$ .


Indoxacarb concentrations ( $\mu\text{g L}^{-1}$ )	Males dry weight (mg)	Females dry weight (mg)
0	0.478 $\pm$ 0.012	0.943 $\pm$ 0.049
1	0.502 $\pm$ 0.014	0.998 $\pm$ 0.030
2	0.461 $\pm$ 0.018	0.981 $\pm$ 0.051
4	0.503 $\pm$ 0.022	0.999 $\pm$ 0.039
8	0.469 $\pm$ 0.012	0.954 $\pm$ 0.049



**Fig. 3.** Oxidative damage, biotransformation, energetic metabolism and neuronal biomarkers in *Chironomus riparius* larvae after 48 h exposure to indoxacarb: a) Lipid Peroxidation; b) DNA Damage; c) Glutathione-S-Transferase; d) Lactate Dehydrogenase; e) Electron Transport System; f) Acetylcholinesterase. All values are presented as mean  $\pm$  SEM,  $n = 7$ . An asterisk denotes statistically significant differences to the control treatment ( $0 \mu\text{g L}^{-1}$ ;  $p < 0.05$ , ANOVA, Dunnett's test).

of  $8 \mu\text{g L}^{-1}$  for larval growth and emergence were observed under exposure to spinosad. Information on the chronic risk to aquatic organisms, including sediment dwellers is lacking (EFSA et al., 2018a). Regarding other dipterans, a spinosad concentration of  $17 \mu\text{g L}^{-1}$  has been

demonstrated to decrease the emergence of *Polypedium nubifer* (Duchet et al., 2015). Concentrations ranging from  $3.7$  to  $45 \mu\text{g L}^{-1}$  seem to affect *Culex pipiens* emergence (Hertlein et al., 2010), while at  $60 \mu\text{g L}^{-1}$  Cetin et al. (2005) reported a complete inhibition of *Culex*



**Fig. 4.** Oxidative stress biomarkers in *Chironomus riparius* larvae after 48 h exposure to indoxacarb: a) Catalase; b) Glutathione Peroxidase; c) Glutathione Reductase; d) Superoxide Dismutase. All values are presented as mean  $\pm$  SEM,  $n = 7$ . An asterisk denotes statistically significant differences to the control treatment ( $0 \mu\text{g L}^{-1}$ ;  $p < 0.05$ , ANOVA, Dunnett's test).

*pipiens* adult emergence. Tomé et al. (2014) determined that exposure to spinosad compromises swimming behavior of *Aedes aegypti*. Behavioral changes have been demonstrated for many neurotoxic compounds and can lead to a reduction in food intake (Pestana et al., 2009, 2010; Tomé et al., 2014; Werner and Moran, 2009), which, although not addressed, might have also occurred here with *C. riparius* and contributed to the reduced growth and developmental rates. Considering other aquatic invertebrates, impairment of population growth rate by spinosad was described for *Daphnia pulex* and *Daphnia magna* at  $8 \mu\text{g L}^{-1}$  (Duchet et al., 2010) and for *Ceriodaphnia dubia* at  $1 \mu\text{g L}^{-1}$  (Deardorff and Stark, 2011).

Regarding the long-term effects of indoxacarb on chironomids, the information available is very limited. Still, a 28-day EC10 of  $1.68 \mu\text{g L}^{-1}$  (endpoint not specified) and a 28-day NOEC (development rate) of  $1.8 \mu\text{g L}^{-1}$  (active substance) were previously determined for *C. riparius* (EFSA et al., 2018b). In the present study, a NOEC of  $4 \mu\text{g L}^{-1}$  was observed for development and emergence endpoints. Ding et al. (2011) investigated the effects of pesticide-contaminated sediments on *C. dilutus*, and the authors concluded that indoxacarb was amongst the most toxic sediment-associated pesticides to *C. dilutus* (10-day LC50 of  $11.3 \mu\text{g g}_{\text{oc}}^{-1}$ ; growth NOEC of  $3.2 \mu\text{g g}_{\text{oc}}^{-1}$ ). Available information shows that commercial formulations of indoxacarb affect life history traits of some insect pest species (Gamil et al., 2011; Martin et al., 2006; Saryazdi et al., 2012), however present results clearly show that life history of non-target aquatic insects may also be altered.

Although the biomarkers exposure experiment does not reflect the entire exposure duration of the chronic toxicity test, short exposures to low concentrations of both insecticides induced several biochemical changes in *C. riparius* larvae. As expected, due to their distinct modes of action, different responses were observed at the biochemical level.

GPx, CAT, and SOD are first-line defense antioxidant enzymes against reactive oxygen species. SOD catalyzes the conversion of superoxide anions to hydrogen peroxide ( $\text{H}_2\text{O}_2$ ), which is subsequently detoxified by CAT and GPx (Ighodaro and Akinloye, 2018). The increase in GPx activity induced by spinosad exposure may have occurred to prevent the accumulation of  $\text{H}_2\text{O}_2$  due to increased oxygen metabolism. GPx has a higher affinity for  $\text{H}_2\text{O}_2$  than CAT (Lushchak, 2012), which may explain why GPx activity increased while catalase activity remained unchanged. The increase in GPx activity was, however, insufficient to prevent oxidative damage, as indicated by the increase of LPO levels and the perceptible increase of DNA damage. The concomitant increase in LPO and GPx has been previously observed in the kidney of *Oreochromis niloticus* (Piner and Uner, 2014) and in mammalian cell lines (Pérez-Perotejo et al., 2008) exposed to the same insecticide. Increased LPO levels are indicative of cellular damage, which may have contributed to the effects observed at the organism level.

Spinosad also led to the increase in ETS activity, an indicator of cellular oxygen metabolism, and LDH activity, involved in the anaerobic pathway of energy production, indicating high levels of energy consumption and high metabolic demand (Rodrigues et al., 2015a; Silva et al., 2016). This increase in energy demand may be associated with the activation of antioxidant mechanisms, as implicit by the increase of GPx and/or other defense mechanisms that were not addressed here. Moreover, an increase in metabolic costs due to the induction of defense mechanisms may also, in part, explain observed reductions in growth and development, since less energy will be available for other physiological processes.

Spinosad's inhibitory effects on AChE activity have been reported for other insect species (El-Mageed and Elgohary, 2006; Maiza et al., 2013; Rabea et al., 2010; Tine et al., 2015), and as a nicotinic acetylcholine receptor modulator, some alterations in AChE activity were anticipated. However, the 48 h exposure to the tested concentrations did not induce changes in AChE activity of *C. riparius* larvae. Azevedo-Pereira et al. (2011) work with *C. riparius* larvae has also revealed that a 48 h exposure to imidacloprid, an insecticide that also targets nicotinic acetylcholine receptors, did not induce alterations in AChE activity. The authors

indicated that inhibitory effects of imidacloprid on AChE were only detected after 96 h of exposure and in the post-exposure period, and yet behavioral changes were linked to AChE activity (Azevedo-Pereira et al., 2011). Given the information available in the literature, it is possible that 48 h exposure to spinosad was not enough to impair AChE activity. Follow-up tests should be performed with prolonged exposure periods, to evaluate the possible extent of spinosad toxic effect on *C. riparius* AChE. Nonetheless, this short-exposure triggered alterations on other biochemical biomarkers, indicating that secondary mechanisms might also be accountable for spinosad's toxicity to *C. riparius*, such as the interference with gamma-aminobutyric acid receptors or others (Salgado and Sparks, 2005).

Regarding the effects of indoxacarb at the biochemical level, GST was the most sensitive endpoint. GST, an enzyme involved in biotransformation and detoxification (Clark, 1989), has been categorized as an ineffective biomarker of pesticide exposure in *C. riparius* (Hirthe et al., 2001), and some works endorse this assumption due to its disparate responses to different pesticides (Planelló et al., 2013). Regardless, in this study, an increase in GST activity as a result of indoxacarb exposure was observed in *C. riparius* larvae. An identical response to indoxacarb was observed in *Blattella germanica* and *Spodoptera littoralis* larvae (Gamil et al., 2011; Maiza et al., 2013). Additionally, Nehare et al. (2010) and Pang et al. (2012) postulated that the detoxification by GST might play a relevant role in indoxacarb resistance. GPx activity also increased in larvae exposed to indoxacarb. Since there were no changes in oxidative damage indicators (LPO and DNA damage), it is suggested that GPx activity and detoxification by GST contributed to preventing oxidative damage in a short-term exposure. As opposed to spinosad, only the anaerobic metabolism (LDH) was induced by indoxacarb in *C. riparius* larvae, since no changes were detected in ETS activity. This induction of LDH may occur due to higher and more readily available energy demands for the activation of GPx and GST, and again this might have contributed to the effects observed at the individual level (reduction in larval growth and increase in time to emergence).

This study elucidates some biochemical responses to spinosad and indoxacarb exposure that may be associated with the effects observed at the organismal level. The induction of defense mechanisms and higher energy expenditures are most likely direct responses of *C. riparius* larvae to cope with the exposure, while oxidative damage may be a direct consequence of spinosad's mechanism of action and may have contributed to the slightly more severe effects observed. Although not specific for insecticide exposure, biochemical biomarkers addressed in the present study may be valuable early-warning tools in biomonitoring studies, since changes were observed in short exposures and at lower concentrations than the ones causing effects at the organismal level. ETS was the most sensitive biochemical biomarker for spinosad, as it was responsive to  $0.5 \mu\text{g L}^{-1}$ , while for indoxacarb GST was the most sensitive biomarker (LOEC of  $4 \mu\text{g L}^{-1}$ ), underlining the role of GST in the detoxification of indoxacarb.

Our findings revealed that under controlled laboratory conditions, spinosad is slightly more toxic to *C. riparius* than indoxacarb since, besides the reduction of larval growth and the increase in time to emergence, a reduction in emergence rate was also observed. Chironomids larvae play a vital role in freshwater ecosystems due to their abundance and food chain position (Péry et al., 2002), and therefore the application of spinosad and indoxacarb near freshwater systems at current rates should be monitored and reviewed, since the concentrations used in this work and that elicited clear deleterious effects are within the estimated environmental levels. Moreover, through their emergence, chironomids represent an important food source for riparian predators, and therefore changes in the emergence of aquatic insects can also have implications for the terrestrial food webs (Schulz et al., 2015).

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.scitotenv.2019.02.303>.

## Acknowledgements

This study had the support of the Fundação para a Ciência e a Tecnologia (FCT) through MARE: UID/MAR/04292/2019; CESAM: UID/AMB/50017/2019, FCT/MCTES through national funds, the co-funding by the FEDER within the PT2020 Partnership Agreement and Compete 2020; Project PROTEOME (PTDC/AAG-MAA/1302/2014) co-financed by COMPETE (POCI-01-0145-FEDER-016773). The project was also partially funded by the Integrated Programme of SR&TD “SmartBioR” (reference Centro-01-0145-FEDER-000018) co-funded by Centro 2020 program, Portugal2020, European Union, through the European Regional Development Fund. Hugo R. Monteiro, Sara C. Novais, and João L.T. Pestana wish to acknowledge the financial support given by FCT (SFRH/BD/80988/2011, SFRH/BPD/94500/2013, IF/01420/2015, respectively).

## References

- Aloísio Torres, M., Pires Testa, C., Gáspari, C., Beatriz Masutti, M., Maria Neves Panitz, C., Curi-Pedrosa, R., Alves de Almeida, E., Di Mascio, P., Wilhelm Filho, D., 2002. Oxidative stress in the mussel *Mytella guyanensis* from polluted mangroves on Santa Catarina Island, Brazil. *Mar. Pollut. Bull.* 44, 923–932. [https://doi.org/10.1016/S0025-326X\(02\)00142-X](https://doi.org/10.1016/S0025-326X(02)00142-X).
- Anikwe, J.C., Adetoro, F.A., Anogwih, J.A., Makanjuola, W.A., Kemabonta, K.A., Akinwande, K.L., 2014. Laboratory and field evaluation of an indoxacarb gel bait against two cockroach species (Dictyoptera: Blattellidae, Blattidae) in Lagos, Nigeria. *J. Econ. Entomol.* 107, 1639–1642. <https://doi.org/10.1603/EC13457>.
- Azevedo-Pereira, H.M.V.S., Lemos, M.F.L., Soares, A.M.V.M., 2011. Effects of imidacloprid exposure on *Chironomus riparius* Meigen larvae: Linking acetylcholinesterase activity to behaviour. *Ecotoxicol. Environ. Saf.* 74, 1210–1215. <https://doi.org/10.1016/j.ecoenv.2011.03.018>.
- Bacci, L., Lupi, D., Savoldelli, S., Rossaro, B., 2016. A review of Spinosyns, a derivative of biological acting substances as a class of insecticides with a broad range of action against many insect pests. *J. Entomol. Acarol. Res.* 48, 40–52. <https://doi.org/10.4081/jea.2016.5653>.
- Bird, R.P., Draper, H.H., 1984. Comparative studies on different methods of malonaldehyde determination. *Methods Enzymol.* 105, 299–305. [https://doi.org/10.1016/S0076-6879\(84\)05038-2](https://doi.org/10.1016/S0076-6879(84)05038-2).
- Bond, J.G., Marina, C.F., Williams, T., 2004. The naturally derived insecticide spinosad is highly toxic to *Aedes* and *Anopheles* mosquito larvae. *Med. Vet. Entomol.* 18, 50–56. <https://doi.org/10.1111/j.0269-283X.2004.0480.x>.
- Boucher, J., Ashley, R., 1999. Effect of spinosad on bell peppers pests and beneficial arthropods. *1998. Arthropod Manage. Tests* 24, E56. <https://doi.org/10.1093/amt/24.1.E56>.
- Cerejeira, M.J., Viana, P., Batista, S., Pereira, T., Silva, E., Valério, M.J., Silva, A., Ferreira, M., Silva-Fernandes, A.M., 2003. Pesticides in Portuguese surface and ground waters. *Water Res.* 37, 1055–1063. [https://doi.org/10.1016/S0043-1354\(01\)00462-6](https://doi.org/10.1016/S0043-1354(01)00462-6).
- Cetin, H., Yanikoglu, A., Cilek, J.E., 2005. Evaluation of the naturally-derived insecticide spinosad against *Culex pipiens* L. (Diptera: Culicidae) larvae in septic tank water in Antalya, Turkey. *J. Vector Ecol.* 30, 151–154.
- Chitgar, M.G., Ghadamyari, M., 2012. Effects of Amitraz on the parasitoid *Encarsia formosa* (Gahan) (Hymenoptera: Aphelinidae) for control of *Trialetodes vaporariorum* Westwood (Homoptera: Aleyrodidae): IOBC methods. *J. Entomol. Res. Soc.* 14 (2), 61–69.
- Clairborne, A., 1985. Catalase activity. In: Greenwald, R.A.E. (Ed.), *Handbook of Methods for Oxygen Radical Research*. CRC Press, Boca Raton, USA, pp. 283–284.
- Clark, A.G., 1989. The comparative enzymology of the glutathione S-transferases from non-vertebrate organisms. *Comp. Biochem. Physiol. B* 92, 419–446. [https://doi.org/10.1016/0305-0491\(89\)90114-4](https://doi.org/10.1016/0305-0491(89)90114-4).
- Cleveland, C.B., Borrett, G.A., Saunders, D.G., Powers, F.L., McGibbon, A.S., Reeves, G.L., Rutherford, L., Balcer, J.L., 2002. Environmental fate of spinosad. 1. Dissipation and degradation in aqueous systems. *J. Agric. Food Chem.* 50, 3244–3256. <https://doi.org/10.1021/jf011663i>.
- Copping, L.G., Menn, J.J., 2000. Biopesticides: a review of their action, applications and efficacy. *Pest Manag. Sci.* 56, 636–676. [https://doi.org/10.1002/1526-4998\(200008\)56:8<651::AID-PS201>3.0.CO;2-U](https://doi.org/10.1002/1526-4998(200008)56:8<651::AID-PS201>3.0.CO;2-U).
- Cribb, A.E., Leeder, J.S., Spielberg, S.P., 1989. Use of a microplate reader in an assay of glutathione reductase using 5,5'-dithiobis(2-nitrobenzoic acid). *Anal. Biochem.* 183, 195–196. [https://doi.org/10.1016/0003-2697\(89\)90188-7](https://doi.org/10.1016/0003-2697(89)90188-7).
- Crouse, G.D., Sparks, T.C., Schoonover, J., Gifford, J., Dripps, J., Bruce, T., Larson, L.L., Garlich, J., Hatton, C., Hill, R.L., Worden, T.V., Martynow, J.G., 2001. Recent advances in the chemistry of spinosyns. *Pest Manag. Sci.* 57, 177–185. [https://doi.org/10.1002/1526-4998\(200102\)57:2<177::AID-PS281>3.0.CO;2-Z](https://doi.org/10.1002/1526-4998(200102)57:2<177::AID-PS281>3.0.CO;2-Z).
- De Coen, W.M., Janssen, C.R., 1997. The use of biomarkers in *Daphnia magna* toxicity testing. IV. Cellular energy allocation: a new methodology to assess the energy budget of toxicant-stressed *Daphnia* populations. *J. Aquat. Ecosyst. Stress. Recover.* 6, 43–55. <https://doi.org/10.1023/A:1008228517955>.
- Deardorff, A., Stark, J., 2011. Population-level toxicity of the insecticide, spinosad and the nonylphenol polyethoxylate, R-11, to the cladoceran species *Ceriodaphnia dubia* Richard. *J. Environ. Sci. Health B* 46, 336–340. <https://doi.org/10.1080/03601234.2011.559891>.
- Diamantino, T.C., Almeida, E., Soares, A.M., Guilhermino, L., 2001. Lactate dehydrogenase activity as an effect criterion in toxicity tests with *Daphnia magna* straus. *Chemosphere* 45, 553–560. [https://doi.org/10.1016/S0045-6535\(01\)00029-7](https://doi.org/10.1016/S0045-6535(01)00029-7).
- Dias, J.L., 2006. *Environmental Fate of Indoxacarb*. California Department of Pesticide Regulation.
- Ding, Y., Weston, D.P., You, J., Rothert, A.K., Lydy, M.J., 2011. Toxicity of sediment-associated pesticides to *Chironomus dilutus* and *Hyalella azteca*. *Arch. Environ. Contam. Toxicol.* 61, 83–92. <https://doi.org/10.1007/s00244-010-9614-2>.
- Dryden, M.W., Payne, P.A., Smith, V., Heaney, K., Sun, F., 2013. Efficacy of indoxacarb applied to cats against the adult cat flea, *Ctenocephalides felis*, flea eggs and adult flea emergence. *Parasit. Vectors* 6, 126. <https://doi.org/10.1186/1756-3305-6-126>.
- Duchet, C., Coutellec, M.-A., Franquet, E., Lagneau, C., Lagadic, L., 2010. Population-level effects of spinosad and *Bacillus thuringiensis israelensis* in *Daphnia pulex* and *Daphnia magna*: comparison of laboratory and field microcosm exposure conditions. *Ecotoxicology* 19, 1224–1237. <https://doi.org/10.1007/s10646-010-0507-y>.
- Duchet, C., Franquet, E., Lagadic, L., Lagneau, C., 2015. Effects of *Bacillus thuringiensis israelensis* and spinosad on adult emergence of the non-biting midges *Polypedium nubifer* (Skuse) and *Tanytarsus curticornis* Kieffer (Diptera: Chironomidae) in coastal wetlands. *Ecotoxicol. Environ. Saf.* 115, 272–278. <https://doi.org/10.1016/j.ecoenv.2015.02.029>.
- EFSA, Arena, M., Auteri, D., Barmaz, S., Bellisai, G., Brancato, A., Brocca, D., Bura, L., Byers, H., Chiusolo, A., Court Marques, D., Crivellente, F., De Lentdecker, C., Egsmose, M., Erdos, Z., Fait, G., Ferreira, L., Goumenou, M., Greco, L., Ippolito, A., Istace, F., Jarrah, S., Kardassi, D., Leuschner, R., Lythgo, C., Magrans, J.O., Medina, P., Miron, I., Molnar, T., Nougadere, A., Padovani, L., Parra Morte, J.M., Pedersen, R., Reich, H., Sacchi, A., Santos, M., Serafimova, R., Sharp, R., Stanek, A., Streissl, F., Sturma, J., Szentes, C., Tarazona, J., Terron, A., Theobald, A., Vagenende, B., Verani, A., Villamar-Bouza, L., 2018a. Peer review of the pesticide risk assessment of the active substance indoxacarb. *Efsa J.* 16 (1), 5140. <https://doi.org/10.2903/j.efsa.2018.5140>.
- EFSA, Arena, M., Auteri, D., Barmaz, S., Brancato, A., Brocca, D., Bura, L., Carrasco Cabrera, L., Chiusolo, A., Court Marques, D., Crivellente, F., De Lentdecker, C., Egsmose, M., Fait, G., Ferreira, L., Goumenou, M., Greco, L., Ippolito, A., Istace, F., Jarrah, S., Kardassi, D., Leuschner, R., Lythgo, C., Magrans, J.O., Medina, P., Miron, I., Molnar, T., Nougadere, A., Padovani, L., Parra Morte, J.M., Pedersen, R., Reich, H., Sacchi, A., Santos, M., Serafimova, R., Sharp, R., Stanek, A., Streissl, F., Sturma, J., Szentes, C., Tarazona, J., Terron, A., Theobald, A., Vagenende, B., Villamar Bouza, L., 2018b. Peer review of the pesticide risk assessment of the active substance spinosad. *Efsa J.* 16 (5), 5252. <https://doi.org/10.2903/j.efsa.2018.5252>.
- Ellman, G.L., Courtney, K.D., Andres, V., Featherstone, R.M., 1961. A new and rapid colorimetric determination of acetylcholinesterase activity. *Biochem. Pharmacol.* 7, 88–95. [https://doi.org/10.1016/0006-2952\(61\)90145-9](https://doi.org/10.1016/0006-2952(61)90145-9).
- El-Mageed, A., Elgohary, L., 2006. Impact of spinosad on some enzymatic activities of the cotton leafworm. *Pak. J. Biol. Sci.* <https://doi.org/10.3923/pjbs.2006.713.716>.
- EPA, 2003. *Indoxacarb; time-limited pesticide tolerance*. *Fed. Regist.* 68, 25824–25831.
- European Commission, 2006. *Commission Directive 2006/10/EC*. *Off. J. Eur. Union* L25, 24–27.
- European Commission, 2008a. *Commission Regulation (Ec) No 404/2008*. *Off. J. Eur. Union* L120, 8–10.
- European Commission, 2008b. *Commission Regulation (Ec) No 889/2008*. *Off. J. Eur. Union* L250, 1–84.
- Gamil, W.E., Mariy, F.M., Youssef, L.A., Abdel Halim, S.M., 2011. Effect of Indoxacarb on some biological and biochemical aspects of *Spodoptera littoralis* (Boisd.) larvae. *Ann. Agric. Sci.* 56, 121–126. <https://doi.org/10.1016/j.aaos.2011.07.005>.
- Guilhermino, L., Lopes, M.C., Carvalho, A.P., Soares, A.M.V.M., 1996. Acetylcholinesterase activity in juveniles of *Daphnia magna* Straus. *Bull. Environ. Contam. Toxicol.* 57, 979–985. <https://doi.org/10.1007/s001289900286>.
- Habig, W.H., Pabst, M.J., Jakoby, W.B., 1974. Glutathione S-transferases. The first enzymatic step in mercapturic acid formation. *J. Biol. Chem.* 249, 7130–7139.
- Hertlein, M.B., Mavrotas, C., Jousseau, C., Lysandrou, M., Thompson, G.D., Jany, W., Ritchie, S.A., 2010. A review of spinosad as a natural product for larval mosquito control. *J. Am. Mosq. Control Assoc.* 26, 67–87. <https://doi.org/10.2987/09-5936.1>.
- Hirth, G., Fisher, T.C., Crane, M., Callaghan, A., 2001. Short-term exposure to sub-lethal doses of lindane affects developmental parameters in *Chironomus riparius* Meigen, but has no effect on larval glutathione-S-transferase activity. *Chemosphere* 44, 583–589. [https://doi.org/10.1016/S0045-6535\(00\)00471-9](https://doi.org/10.1016/S0045-6535(00)00471-9).
- Ighodaro, O.M., Akinloye, O.A., 2018. First line defence antioxidants-superoxide dismutase (SOD), catalase (CAT) and glutathione peroxidase (GPx): their fundamental role in the entire antioxidant defence grid. *Alexandria J. Med.* 54, 287–293.
- Jones, T., Scott-Dupree, C., Harris, R., Shipp, L., Harris, B., 2005. The efficacy of spinosad against the western flower thrips, *Frankliniella occidentalis*, and its impact on associated biological control agents on greenhouse cucumbers in southern Ontario. *Pest Manag. Sci.* 61, 179–185. <https://doi.org/10.1002/ps.939>.
- de Lafontaine, Y., Gagné, F., Blaise, C., Costan, G., Gagnon, P., Chan, H.M., 2000. Biomarkers in zebra mussels (*Dreissena polymorpha*) for the assessment and monitoring of water quality of the St Lawrence River (Canada). *Aquat. Toxicol.* 50, 51–71. [https://doi.org/10.1016/S0166-445X\(99\)00094-6](https://doi.org/10.1016/S0166-445X(99)00094-6).
- Lahm, G.P., McCann, S.F., Harrison, C.R., Stevenson, T.M., Shapiro, R., 2000. Evolution of the sodium channel blocking insecticides: the discovery of indoxacarb. *Agrochemical Discovery, Insect, Weed, and Fungal Control*. American Chemical Society, Washington, DC, pp. 20–34. <https://doi.org/10.1021/bk-2001-0774.ch003>.
- Lapied, B., Grolleau, F., Sattelle, D.B., 2001. Indoxacarb, an oxadiazine insecticide, blocks insect neuronal sodium channels. *Br. J. Pharmacol.* 132, 587–595. <https://doi.org/10.1038/sj.bjp.0703853>.
- Lawler, S.P., Dritz, D.A., 2013. Efficacy of spinosad in control of larval *Culex tarsalis* and chironomid midges, and its nontarget effects. *J. Am. Mosq. Control Assoc.* 29, 352–357. <https://doi.org/10.2987/13-6369.1>.
- Lemos, M.F.L., Soares, A.M.V.M., Correia, A.C., Esteves, A.C., 2010. Proteins in ecotoxicology - how, why and why not? *Proteomics* 10, 873–887. <https://doi.org/10.1002/pmic.200900470>.

- Liu, T.-X., Zhang, Y., 2012. Side effects of two reduced-risk insecticides, indoxacarb and spinosad, on two species of *Trichogramma* (Hymenoptera: Trichogrammatidae) on cabbage. *Ecotoxicology* 21, 2254–2263. <https://doi.org/10.1007/s10646-012-0981-5>.
- Lushchak, V.I., 2012. Glutathione homeostasis and functions: potential targets for medical interventions. *J. Amino Acids* 2012, 1–26. <https://doi.org/10.1155/2012/736837>.
- Maiza, A., Aribi, N., Smaghe, G., Kilani-Morakchi, S., Bendjedid, M., Soltani, N., 2013. Sub-lethal effects on reproduction and survival by spinosad and indoxacarb in cockroaches *Blattella Germanica*. *Bull. Insectology* 66 (1), 11–20.
- Martin, N.A., Workman, P.J., Hedderley, D., 2006. Susceptibility of *Scaptomyza flava* (Diptera: Drosophilidae) to insecticides. *N. Z. Plant Prot.* 59, 228–234.
- McCord, J.M., Fridovich, I., 1969. Superoxide dismutase. An enzymic function for erythrocyte (hemocuprein). *J. Biol. Chem.* 244, 6049–6055.
- Mertz, F.P., Yao, R.C., 1990. *Saccharopolyspora spinosa* sp. nov. isolated from soil collected in a sugar mill rum still. *Int. J. Syst. Bacteriol.* 40, 34–39. <https://doi.org/10.1099/0020713-40-1-34>.
- Mohandas, J., Marshall, J.J., Duggin, G.G., Horvath, J.S., Tiller, D.J., 1984. Differential distribution of glutathione and glutathione-related enzymes in rabbit kidney: Possible implications in analgesic nephropathy. *Biochem. Pharmacol.* 33, 1801–1807. [https://doi.org/10.1016/0006-2952\(84\)90353-8](https://doi.org/10.1016/0006-2952(84)90353-8).
- Monteiro, H.R., Lemos, M.F.L., Novais, S.C., Soares, A.M.V.M., Pestana, J.L.T., 2019. Amitraz toxicity to the midge *Chironomus riparius*: life-history and biochemical responses. *Chemosphere* 221, 324–332. <https://doi.org/10.1016/j.chemosphere.2019.01.018>.
- Nehare, S., Moharil, M.P., Ghodki, B.S., Lande, G.K., Bisane, K.D., Thakare, A.S., Barkhade, U.P., 2010. Biochemical analysis and synergistic suppression of indoxacarb resistance in *Plutella xylostella* L. *J. Asia Pac. Entomol.* 13, 91–95. <https://doi.org/10.1016/j.aspen.2009.12.002>.
- OECD, 2004. Test no. 219: sediment–water chironomid toxicity using spiked water. OECD Guidel Test Chem. OECD Publishing. <https://doi.org/10.1787/9789264070288-en>.
- OECD, 2011. Test no. 235: *Chironomus* sp., acute immobilisation test. OECD Guidel Test Chem. OECD Publishing. <https://doi.org/10.1787/9789264122383-en>.
- Ohkawa, H., Ohishi, N., Yagi, K., 1979. Assay for lipid peroxides in animal tissues by thiobarbituric acid reaction. *Anal. Biochem.* 95, 351–358. [https://doi.org/10.1016/0003-2697\(79\)90738-3](https://doi.org/10.1016/0003-2697(79)90738-3).
- Olive, P.L., 1988. DNA precipitation assay: A rapid and simple method for detecting DNA damage in mammalian cells. *Environ. Mol. Mutagen.* 11, 487–495. <https://doi.org/10.1002/em.2850110409>.
- Orr, N., Shaffner, A.J., Richey, K., Crouse, G.D., 2009. Novel mode of action of spinosad: receptor binding studies demonstrating lack of interaction with known insecticidal target sites. *Pestic. Biochem. Physiol.* 95, 1–5. <https://doi.org/10.1016/j.pestbp.2009.04.009>.
- Oxborough, R.M., N'Guessan, R., Kitau, J., Tungu, P.K., Malone, D., Mosha, F.W., Rowland, M.W., 2015. A new class of insecticide for malaria vector control: evaluation of mosquito nets treated singly with indoxacarb (oxadiazine) or with a pyrethroid mixture against *Anopheles gambiae* and *Culex quinquefasciatus*. *Malar. J.* 14. <https://doi.org/10.1186/s12936-015-0890-1>.
- Pang, S., You, W., Duan, L., Song, X., Li, X., Wang, C., 2012. Resistance selection and mechanisms of oriental tobacco budworm (*Helicoverpa assulta* Guenee) to indoxacarb. *Pestic. Biochem. Physiol.* 103, 219–223. <https://doi.org/10.1016/j.pestbp.2012.05.011>.
- Pérez, C.M., Marina, C.F., Bond, J.G., Rojas, J.C., Valle, J., Williams, T., 2007. Spinosad, a naturally derived insecticide, for control of *Aedes aegypti* (Diptera: Culicidae): efficacy, persistence, and elicited oviposition response. *J. Med. Entomol.* 44, 631–638. <https://doi.org/10.1093/jmedent/44.4.631>.
- Pérez-Pertejo, Y., Reguera, R.M., Ordóñez, D., Balaña-Fouce, R., 2008. Alterations in the glutathione–redox balance induced by the bio-insecticide Spinosad in CHO-K1 and Vero cells. *Ecotoxicol. Environ. Saf.* 70, 251–258. <https://doi.org/10.1016/j.ecoenv.2007.06.009>.
- Péry, A.R.R., Mons, R., Flammarion, P., Lagadic, L., Garric, J., 2002. A modeling approach to link food availability, growth, emergence, and reproduction for the midge *Chironomus riparius*. *Environ. Toxicol. Chem.* 21, 2507–2513. <https://doi.org/10.1002/etc.5620211133>.
- Pestana, J.L.T., Alexander, A.C., Culp, J.M., Baird, D.J., Cessna, A.J., Soares, A.M.V.M., 2009. Structural and functional responses of benthic invertebrates to imidacloprid in outdoor stream mesocosms. *Environ. Pollut.* 157, 2328–2334. <https://doi.org/10.1016/j.envpol.2009.03.027>.
- Pestana, J.L.T., Loureiro, S., Baird, D.J., Soares, A.M.V.M., 2010. Pesticide exposure and inducible antipredator responses in the zooplankton grazer, *Daphnia magna* Straus. *Chemosphere* 78, 241–248. <https://doi.org/10.1016/j.chemosphere.2009.10.066>.
- Piner, P., Uner, N., 2014. Organic insecticide spinosad causes in vivo oxidative effects in the brain of *Oreochromis niloticus*. *Environ. Toxicol.* 29, 253–260. <https://doi.org/10.1002/tox.21753>.
- Planelló, R., Servia, M.J., Gómez-Sande, P., Herrero, O., Cobo, F., Morcillo, G., 2013. Transcriptional responses, metabolic activity and mouthpart deformities in natural populations of *Chironomus riparius* larvae exposed to environmental pollutants. *Environ. Toxicol.* 30, 383–395. <https://doi.org/10.1002/tox.21893>.
- Pridgeon, J.W., Becnel, J.J., Clark, G.G., Linthicum, K.J., 2009. A high-throughput screening method to identify potential pesticides for mosquito control. *J. Med. Entomol.* 46, 335–341. <https://doi.org/10.1603/033.046.0219>.
- Rabea, E.I., Nasr, H.M., Badawy, M.E.I., 2010. Toxic effect and biochemical study of chlorfluazuron, oxydemeton, and spinosad on honey bees (*Apis mellifera*). *Arch. Environ. Contam. Toxicol.* 58, 722–732. <https://doi.org/10.1007/s00244-009-9403-y>.
- Rodrigues, A.C.M., Gravato, C., Quintaneiro, C., Barata, C., Soares, A.M.V.M., Pestana, J.L.T., 2015a. Sub-lethal toxicity of environmentally relevant concentrations of esfenvalerate to *Chironomus riparius*. *Environ. Pollut.* 207, 273–279. <https://doi.org/10.1016/j.envpol.2015.09.035>.
- Rodrigues, A.C.M., Pestana, J.L.T., 2015b. Life history and biochemical effects of chlorantraniliprole on *Chironomus riparius*. *Sci. Total Environ.* 508, 506–513. <https://doi.org/10.1016/j.scitotenv.2014.12.021>.
- Salgado, V.L., 1998. Studies on the mode of action of spinosad: insect symptoms and physiological correlates. *Pestic. Biochem. Physiol.* 60, 91–102. <https://doi.org/10.1006/pest.1998.2332>.
- Salgado, V.L., Sparks, T.C., 2005. The spinosyns: chemistry, biochemistry, mode of action, and resistance. In: Gilbert, L.I. (Ed.), *Comprehensive Molecular Insect Science*. Elsevier, Amsterdam, Netherlands, pp. 137–173. <https://doi.org/10.1016/B0-44-51924-6/00078-8>.
- Salgado, V.L., Sheets, J.J., Watson, G.B., 1998. Studies on the mode of action of spinosad: the internal effective concentration and the concentration dependence of neural excitation. *Pestic. Biochem. Physiol.* 60, 103–110. <https://doi.org/10.1006/pest.1998.2333>.
- Sarfraz, M., Dossall, L.M., Keddie, B.A., 2005. Spinosad: a promising tool for integrated pest management. *Outlook Pest Man.* 16, 78–84. <https://doi.org/10.1564/16ap109>.
- Saryzadi, A., Ghasem, Hejazi, J. Mir, Saber, M., 2012. Residual toxicity of abamectin, chlorpyrifos, cyromazine, indoxacarb and spinosad on *Liriomyza trifolii* (Burgess) (Diptera: Agromyzidae) in greenhouse conditions. *Pesticidi i fitomedicina* 27, 107–116. <https://doi.org/10.2298/PIF1202107S>.
- Schulz, R., 2004. Field studies on exposure, effects, and risk mitigation of aquatic nonpoint-source insecticide pollution: a review. *J. Environ. Qual.* 33, 419–448. <https://doi.org/10.2134/jeq2004.0419>.
- Schulz, R., Bundschuh, M., Gergs, R., Brühl, C.A., Diehl, D., Entling, M.H., Fahse, L., Frör, O., Jungkunst, H.F., Lorke, A., Schafer, R.B., Schaumann, G.E., Schwenk, K., 2015. Review on environmental alterations propagating from aquatic to terrestrial ecosystems. *Sci. Total Environ.* 538, 246–261. <https://doi.org/10.1016/j.scitotenv.2015.08.038>.
- Sibley, P.K., Benoit, D.A., Ankley, G.T., 1997. The significance of growth in *Chironomus tentans* sediment toxicity tests: relationship to reproduction and demographic endpoints. *Environ. Toxicol. Chem.* 16, 336. <https://doi.org/10.1002/etc.5620160232>.
- Sibley, P.K., Ankley, G.T., Benoit, D.A., 2001. Factors affecting reproduction and the importance of adult size on reproductive output of the midge *Chironomus tentans*. *Environ. Toxicol. Chem.* 20, 1296. <https://doi.org/10.1002/etc.5620200618>.
- Silva, C.S.E., Novais, S.C., Lemos, M.F.L., Mendes, S., Oliveira, A.P., Gonçalves, E.J., Faria, A.M., 2016. Effects of ocean acidification on the swimming ability, development and biochemical responses of sand smelt larvae. *Sci. Total Environ.* 563–564, 89–98. <https://doi.org/10.1016/j.scitotenv.2016.04.091>.
- Stara, J., Ourednickova, J., Kocourek, F., 2010. Laboratory evaluation of the side effects of insecticides on *Aphidius colemani* (Hymenoptera: Aphididae), *Aphidoletes aphidimyza* (Diptera: Cecidomyiidae), and *Neoseiulus cucumeris* (Acari: Phytoseiidae). *J. Pest. Sci.* 84, 25–31. <https://doi.org/10.1007/s10340-010-0322-5>.
- Stevens, M.M., Helliwell, S., Hughes, P.A., 2005. Toxicity of *Bacillus thuringiensis* var. *israelensis* formulations, spinosad, and selected synthetic insecticides to *Chironomus tepperi* larvae. *J. Am. Mosq. Control Assoc.* 21, 446–450. [https://doi.org/10.2987/8756-971X\(2006\)21\[446:TOBTVI\]2.0.CO;2](https://doi.org/10.2987/8756-971X(2006)21[446:TOBTVI]2.0.CO;2).
- Thompson, G.D., Dutton, R., Sparks, T.C., 2000. Spinosad – a case study: an example from a natural products discovery programme. *Pest Manag. Sci.* 56, 696–702. [https://doi.org/10.1002/1526-4998\(200008\)56:8<696::AID-PS182>3.0.CO;2-5](https://doi.org/10.1002/1526-4998(200008)56:8<696::AID-PS182>3.0.CO;2-5).
- Tine, S., Tine-Djebbar, F., Aribi, N., Boudjelida, H., 2015. Topical toxicity of spinosad and its impact on the enzymatic activities and reproduction in the cockroach *Blattella orientalis* (Dictyoptera: Blattellidae). *Afr. Entomol.* 23, 387–396. <https://doi.org/10.4001/003.023.0230>.
- Tomé, H.V., Pascini, T.V., Dângelo, R.A., Guedes, R.N., Martins, G.F., 2014. Survival and swimming behavior of insecticide-exposed larvae and pupae of the yellow fever mosquito *Aedes aegypti*. *Parasit. Vectors* 7, 195. <https://doi.org/10.1186/1756-3305-7-195>.
- Vassault, A., 1983. Lactate dehydrogenase. In: Bergmeyer, H.U., Bergmeyer, J., Grafl, M. (Eds.), *Methods of Enzymatic Analysis*, third ed. vol. III. Weinheim, Verlag Chemie, pp. 118–126.
- Weltje, L., Rufli, H., Heimbach, F., Wheeler, J., Vervliet-Scheebaum, M., Hamer, M., 2010. The chironomid acute toxicity test: development of a new test system. *Integr. Environ. Assess. Manag.* 6, 301–307. [https://doi.org/10.1897/IEAM\\_2009-069.1](https://doi.org/10.1897/IEAM_2009-069.1).
- Werner, I., Moran, K., 2009. Effects of pyrethroid insecticides on aquatic organisms. *Agrochemical Discovery, Occurrence and Behavior in Aquatic Environments*. American Chemical Society, Washington, DC, USA, pp. 310–334. <https://doi.org/10.1021/bk-2008-0991.ch014>.
- Wilkinson, J.D., Biever, K.D., Ignoffo, C.M., 1979. Synthetic pyrethroid and organophosphate insecticides against the parasitoid *Apanteles marginiventris* and the predators *Geocoris punctipes*, *Hippodamia convergens*, and *Podisus maculiventris*. *J. Econ. Entomol.* 72, 473–475. <https://doi.org/10.1093/jee/72.4.473>.
- Wing, K.D., Schnee, M.E., Sacher, M., Connair, M., 1998. A novel oxadiazine insecticide is bioactivated in lepidopteran larvae. *Arch. Pharm. Res.* 37, pp. 91–10391. [https://doi.org/10.1002/\(SICI\)1520-6327\(1998\)37:1<91::AID-ARCH11>3.0.CO;2-5](https://doi.org/10.1002/(SICI)1520-6327(1998)37:1<91::AID-ARCH11>3.0.CO;2-5).
- Wing, K.D., Sacher, M., Kagaya, Y., Tsurubuchi, Y., Mulderig, L., Connair, M., Schnee, M., 2000. Bioactivation and mode of action of the oxadiazine indoxacarb in insects. *Crop Prot.* 19, 537–545. [https://doi.org/10.1016/S0261-2194\(00\)00070-3](https://doi.org/10.1016/S0261-2194(00)00070-3).